

COMBINATORICS AND TOPOLOGY OF STRAIGHTENING MAPS II: DISCONTINUITY

HIROYUKI INOU

ABSTRACT. We continue the study of straightening maps for the family of polynomials of degree $d \geq 3$. The notion of straightening map is originally introduced by Douady and Hubbard to study the self-similarity of the Mandelbrot set. As expected from their example of a cubic-like family with discontinuous straightening map, we prove that the straightening map is discontinuous unless it is of disjoint type (in this case, it is already known that the straightening map is continuous).

1. INTRODUCTION

Consider the family of monic centered polynomials $\text{Poly}(d)$ of degree $d \geq 2$. The *connectedness locus* $\mathcal{C}(d)$ is the set of $f \in \text{Poly}(d)$ having connected filled Julia set $K(f)$. When $d = 2$, $\mathcal{M} = \mathcal{C}(2)$ is the well-known Mandelbrot set. Douady and Hubbard [DH85] proved that there exist infinitely many small copies of the Mandelbrot set in itself. In fact, for any $z^2 + c_0 \in \mathcal{M}$ such that the critical point 0 is periodic, there exists a homeomorphism $\chi : M' \rightarrow \mathcal{M}$ such that $c_0 \in M'$, the boundary of M' is contained in that of $\mathcal{M}$ and $\chi(c_0) = 0$ [Hai00].

The map χ above is an example of what we call a *straightening map*. For a family of polynomial-like mappings parameterized by a complex manifold Λ of degree $d \geq 2$, we can define such a map defined on the connectedness locus of Λ to the set of affine conjugacy classes of polynomials in $\mathcal{C}(d)$.

In the preceding paper [IK08], we consider straightening maps for the family of renormalizable polynomials of degree $d \geq 3$. The combinatorics of a family of renormalizable polynomials can be described in terms of rational lamination introduced by Thurston [Thu85]. A *rational lamination* λ_f for $f \in \mathcal{C}(d)$ is the landing relation of external rays of rational angles. Let λ_0 be a post-critically finite d -invariant rational lamination, and let $\mathcal{C}(\lambda_0) = \{f \in \mathcal{C}(d); \lambda_f \supset \lambda_0\}$ denote the set of λ_0 -combinatorially renormalizable polynomials. For $f \in \mathcal{C}(\lambda_0)$, we say f is λ_0 -renormalizable if it has a polynomial-like restriction whose filled Julia sets are λ_0 -fibers, which are continua defined in terms of λ_0 . Hence we can straighten such a restriction (λ_0 -renormalization) to get a new polynomial by the straightening theorem by Douady and Hubbard [DH85]. More precisely, since there might exist several critical points, we borrow the notion of mapping schema introduced by Milnor [MP92] to describe the dynamics of λ_0 -fibers containing critical points. Therefore, the straightening of a λ_0 -renormalization of f is an affine conjugacy class of polynomials over a mapping schema of λ_0 .

Under this definition, the straightening map is at most finite-to-one. By introducing “markings” for polynomials and polynomial-like maps, we can define an *injective* straightening map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \rightarrow \mathcal{C}(T(\lambda_0))$, where $\mathcal{C}(T(\lambda_0))$ is the *fiberwise connectedness locus*

of the family $\text{Poly}(T(\lambda_0))$ of monic centered polynomials over $T(\lambda_0)$. We recall these notions and results in the preceding paper in Section 6.

In this paper, we study discontinuity of straightening maps.

Main Theorem. *Let $d \geq 3$. If a d -invariant post-critically finite rational lamination λ_0 has nonempty domain and a non-trivial Fatou critical relation, the straightening map χ_{λ_0} is not continuous. More precisely, χ_{λ_0} is not continuous on any neighborhood of any Misiurewicz λ_0 -renormalizable polynomial.*

We say a polynomial f of degree $d \geq 2$ is Misiurewicz if all critical points are (strictly) preperiodic. Note that since Misiurewicz maps are quasiconformally rigid (moreover, they are combinatorially rigid), straightening maps are continuous at those parameters.

It is known that the closure of the set of Misiurewicz polynomials coincides with the support of bifurcation measure [DF08]. Therefore, we may also say that χ_{λ_0} is not continuous on any open set intersecting the support of the bifurcation measure.

An equivalent condition for the domain to be nonempty is stated in [IK08] (see Proposition 6.8).

It is well-known that straightening maps for quadratic-like families are always continuous [DH85]. Therefore, we have a complete classification:

Corollary 1.1. *Let $d \geq 3$ and let λ_0 be a d -invariant post-critically finite rational lamination. Then the straightening map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \rightarrow \mathcal{C}(T(\lambda_0))$ is continuous if and only if λ_0 is of disjoint type.*

We say a d -invariant post-critically finite rational lamination λ_0 is of *disjoint type* if it is the rational lamination of a post-critically finite polynomial such that all Fatou critical points are simple and periodic, and lie in different orbits (an equivalent definition without polynomial realization is given in Section 6). In this case, renormalizations consist of quadratic-like mappings, so the straightening map is continuous.

Also, in the same article, Douady and Hubbard have already given an example of cubic-like family whose straightening map is discontinuous. Although their example strongly suggests that straightening maps are often discontinuous, their example is constructed by putting some invariant complex dilatation outside filled Julia sets of polynomials, and their argument does not allow us to know whether a given straightening map is continuous or not.

Epstein [Eps] have also proved that straightening maps are discontinuous on the boundary of the *main* hyperbolic component (i.e., the one containing the power map), and his result can be generalized to all hyperbolic components such that an attracting periodic orbit attracts at least two critical points, by use of the author's result [Ino08]. Epstein's result and our result have many similarities; both depend on parabolic implosion, and prove existence of an analytic conjugacy between renormalization and its straightening assuming that the straightening map is continuous. However, the proofs for the existence are completely different. Epstein's proof depends on analytic dependence of Ecalle-Voronin invariants and our proof depends on combinatorial constructions with the help of rational laminations. This difference yields completely different sequences which cause discontinuity in the limit; Epstein's one is in a hyperbolic component, and ours is in the bifurcation locus.

In the case of cubic polynomials, fully renormalizable polynomials are divided into four types in terms of mapping schema, according to Milnor [Mil92]; adjacent, bitransitive (bicritical), capture and disjoint. The target space of straightening maps are determined by these types: It is the cubic connectedness locus $\mathcal{C}(3)$ for adjacent type, the connectedness locus of biquadratic family $\mathcal{C}(2 \times 2) = \{(a, b) \in \mathbb{C}^2; K((z^2 + a)^2 + b) \text{ is connected}\}$ for

bitransitive type, the full family of connected quadratic filled Julia sets $\mathcal{MK} = \{(c, z); c \in \mathcal{M}, z \in K(z^2 + c)\}$ for capture type, and the product space of the Mandelbrot set with itself $\mathcal{M} \times \mathcal{M}$ for disjoint type. Any disjoint type straightening map is continuous because it consists of straightening maps of two quadratic-like families. All the other cases have a non-trivial Fatou critical relations, so straightening maps are not continuous.

On the other hand, for capture renormalizations, straightening maps are continuous on each fiber. Buff and Henriksen [BH01] have proved there is a natural quasiconformal embedding of the filled Julia set $K(\lambda z + z^2)$ for $|\lambda| \leq 1$ into the connectedness locus of a cubic one-parameter family, and we proved that any connected filled Julia set can be homeomorphically embedded to the connectedness locus of any higher degree polynomial [Ino06]. Furthermore, we have proved in the preceding paper [IK08] that for a cubic rational lamination of primitive capture type, the straightening map χ is surjective onto $\mathcal{MK}$, and its restriction to $\mathcal{K}_c = \chi^{-1}(\{c\} \times K(z^2 + c))$ for each $c \in \mathcal{M}$ can be extended to a quasiconformal embedding, possibly after desingularizing the one-dimensional analytic set containing $\mathcal{K}_c$. Therefore, by Main Theorem, such quasiconformal embeddings of connected Julia sets does not move continuously on polynomials.

The most of the proof of Main Theorem can be also applied to renormalizable rational maps and transcendental entire maps. We discuss that in the last section (Section 11).

Our argument needs two-dimensional bifurcations to prove discontinuity: One is bifurcation of two critical orbits in one grand orbit, and the other is parabolic bifurcation. Therefore, we cannot apply our argument to a one-parameter family of polynomials. Hence it is natural to ask whether we can get discontinuous straightening maps for smaller parameter spaces. And we may also ask whether parabolic bifurcation is the unique possibility to get discontinuity. Here is a list of those questions:

Question. Can straightening maps be discontinuous under the following conditions?

- (i) On real polynomial families.
- (ii) On dynamically defined complex one-parameter spaces.
- (iii) On anti-holomorphic one-parameter families.
- (iv) At non-parabolic parameters (having a Siegel disk, or an invariant line field on the Julia set).

The first and third parameter spaces can contain “baby tricones”, and they might be nice candidates to study.

The proof of Main Theorem consists of several steps. The first step is to relate the continuity of a straightening map for an analytic family of polynomial-like mappings with two marked points (abbr. AFPL2MP) to the multipliers of repelling periodic orbits (Theorem 5.1). Here we consider a similar situation as the example of discontinuous straightening map by Douady and Hubbard. We start with a polynomial-like map having a parabolic periodic point whose basin contains both of the marked points. If the straightening map is continuous in a neighborhood of this map, and it has nice perturbations described in terms of a given repelling periodic point, parabolic implosion and Lavaurs map, then the modulus of the multiplier of the repelling periodic point does not change by straightening. Two marked points will be critical orbits in the application, so that the continuity of χ_{λ_0} implies the continuity of the straightening map of this AFPL2MP.

Secondly, we study parabolic bifurcations to find nice perturbations so that we can apply the first step (Section 8). Next, we find a nice parabolic map f_1 arbitrarily close to a given Misiurewicz polynomial f_0 so that we can apply the second step (Section 9.1).

By gluing these three steps together, if the straightening map χ is continuous in a neighborhood of f_0 , we can get a hybrid conjugacy preserving multipliers between quadratic-like restrictions of some iterates of a renormalization of f_1 and its straightening $P_1 = \chi(f_1)$. Thus they are analytically conjugate by Sullivan-Prado-Przytycki-Urbanski theorem (Theorem 2.3).

Then, applying the results on analytic conjugate polynomial-like restrictions of polynomials [Ino08] to get a contradiction.

Since this proof is constructive, we can specify at which parameter a straightening map is not continuous. See Remark 10.1 for details.

One of the most difficulties in the proof is that we need to perturb inside the connectedness locus. Moreover, we need to perturb in the domain of the straightening map $\mathcal{R}(\lambda_0)$. To do this, we construct a rational lamination or a critical portrait which the desired map should have, then we realize them by polynomials. However, since those combinatorial objects are not complete invariants, we cannot apply this construction parabolic polynomials. Hence we first construct Misiurewicz polynomials and take a limit to find some of those perturbations. To show that Misiurewicz polynomials constructed in this way and their limits are in $\mathcal{R}(\lambda_0)$, we also need some facts that $\mathcal{R}(\lambda_0)$ contains plenty of dynamics (Theorem 6.9 and Theorem 6.12), proved in the preceding paper.

Acknowledgment. The author would thank Mitsuhiro Shishikura for helpful comments. He would also thank Peter Haïssinsky, Tomoki Kawahira and Jan Kiwi for valuable discussions. He would also like to express his gratitude to Institut de Mathématiques de Toulouse for its hospitality during his visit during 2007/2008 when this paper was written.

2. POLYNOMIAL-LIKE MAPPINGS

Definition (Polynomial-like mapping). A *polynomial-like mapping* is a proper holomorphic map $f : U' \rightarrow U$ with $U' \Subset U \subset \mathbb{C}$. We always assume the degree of f is at least two. The *filled Julia set* $K(f) = K(f; U', U)$ is defined by

$$K(f; U', U) = \bigcap_{n \geq 0} f^{-n}(U')$$

and we call $J(f) = J(f; U', U) = \partial K(f; U', U)$ the *Julia set*.

We introduce the notion of *external markings*, which is necessary to distinguish polynomials whose renormalizations are hybrid equivalent but combinatorially different.

Definition (Access and external marking). Let $f : U' \rightarrow U$ is a polynomial-like mapping. A *path to* $K(f)$ is a path $\gamma : [0, 1] \rightarrow U'$ such that $\gamma(0) \in J(f)$ and $\gamma((0, 1]) \subset U' \setminus K(f)$. For a path γ to $K(f)$, there exists a unique component of $f(\gamma) \cap U'$ which is also a path to $K(f)$ (after a suitable reparametrization). We denote it by $f_*\gamma$.

We say two paths γ_0, γ_1 to $K(f)$ are *homotopic* if they are homotopic rel $K(f)$, i.e., if there exists a homotopy $\gamma : [0, 1] \times [0, 1] \rightarrow U'$ such that $\gamma(0, t) = \gamma_0(t)$, $\gamma(1, t) = \gamma_1(t)$ and $\gamma(s, 0) = \gamma_0(0)$. An *access* for $f : U' \rightarrow U$ is a homotopy class of paths to $K(f)$. We say an access $[\gamma]$ is *invariant* if $f_*\gamma$ is homotopic to γ . It is easy to see that this definition does not depend on the choice of representatives.

An *external marking* of a polynomial-like mapping is an invariant access. An *externally marked polynomial-like mapping* is a pair $(f : U' \rightarrow U, [\gamma])$ of a polynomial-like mapping and an external marking of it.

Example. Let f be a monic centered polynomial of degree $d \geq 2$. For sufficiently large $R > 0$, let $U = \Delta(R) = \{|z| < R\}$ and $U' = f^{-1}(U)$. Then $f : U' \rightarrow U$ is a polynomial-like

mapping of degree d . If the external ray $R_f(0)$ of angle 0 does not bifurcate (e.g., when $K(f)$ is connected), it lands at a fixed point in $J(f)$ and defines an external marking for it. We call it the *standard external marking* for f .

Let $\text{Poly}(d)$ be the family of monic centered polynomials of degree d and let $\mathcal{C}(d)$ be its connectedness locus, i.e., the set of all $f \in \text{Poly}(d)$ such that the filled Julia set $K(f)$ is connected.

By use of the standard external marking, $\mathcal{C}(d)$ can be considered as the set of affine conjugacy classes of externally marked polynomials of degree d with connected Julia sets.

Definition (Hybrid equivalence). We say two polynomial-like mappings $f : U' \rightarrow U$ and $g : V' \rightarrow V$ are *hybrid equivalent* if there exists a quasiconformal homeomorphism $\psi : U'' \rightarrow V''$ between neighborhoods of the filled Julia sets of f and g such that $\psi \circ f = g \circ \psi$ and $\bar{\partial}\psi \equiv 0$ a.e. on $K(f; U', U)$.

For externally marked polynomial-like mappings $(f, [\gamma_f])$ and $(g, [\gamma_g])$, we say a hybrid conjugacy ψ between f and g *respects external markings* if $\psi(\gamma_f)$ is homotopic to γ_g .

The following theorem by Douady and Hubbard [DH85] classifies polynomial-like mappings in the sense of hybrid conjugacy. It also asserts that most dynamical properties for polynomials also holds for polynomial-like mappings. We can further add some information on external markings (see [IK08]).

Theorem 2.1 (Straightening theorem). *Any polynomial-like mapping $f : U' \rightarrow U$ of degree d is hybrid equivalent to some polynomial $g \in \text{Poly}(d)$.*

Moreover, if $K(f; U', U)$ is connected and $f : U' \rightarrow U$ is externally marked, then such a polynomial $g \in \mathcal{C}(d)$ is unique assuming that a hybrid conjugacy respects the external markings, where the external marking of g is the standard external marking.

For a periodic point $x \in \mathbb{C}$ of period n for a holomorphic map f , let us denote its multiplier by $\text{mult}_f(x)$, i.e.,

$$\text{mult}_f(x) = (f^n)'(x).$$

Definition (Hybrid conjugacy preserving multipliers). Let $f : U' \rightarrow U$ and $g : V' \rightarrow V$ are polynomial-like mappings and $\psi : U \rightarrow V$ be a hybrid conjugacy. We say that ψ *preserves multipliers* if for any periodic point x for f , we have

$$|\text{mult}_f(x)| = |\text{mult}_g(\psi(x))|.$$

Definition (Conjugate by an irreducible holomorphic correspondence). We say two rational maps f_1 and f_2 are *conjugate by an irreducible holomorphic correspondence* if there exist rational maps g, ψ_1 and ψ_2 such that $\psi_i \circ g = f_i \circ \psi_i$.

In particular, when f_1 and f_2 are conjugate by an irreducible holomorphic correspondence, they have the same degree.

The aim of this section is to prove the following:

Theorem 2.2. *Let f_1 and f_2 be rational maps. Assume they have polynomial-like restrictions $f_i : U'_i \rightarrow U_i$, $i = 1, 2$ which are hybrid conjugate by a conjugacy preserving multipliers. Then f_1 and f_2 are conjugate by an irreducible holomorphic correspondence.*

The following theorem is essentially proved by Prado [Pra96] based on the idea given by Sullivan [Sul87], and its complete proof was given by Przytycki and Urbanski [PU99].

Theorem 2.3 (Sullivan-Prado-Przytycki-Urbanski). *Suppose that $f : U' \rightarrow U$ and $g : V \rightarrow V$ are two tame polynomial-like maps. Then the following are equivalent:*

- (i) *there exists a hybrid conjugacy between f and g preserving multipliers.*
- (ii) *f and g are analytically conjugate, i.e., there exists a conformal isomorphism $\varphi : U'' \rightarrow V''$ conjugating f and g , where U'' and V'' are neighborhoods of $K(f)$ and $K(g)$ respectively.*

Remark 2.4. Although Przytycki and Urbanski proved the theorem in the case of rational maps, the same proof can be applied for polynomial-like mappings. Moreover, they only proved the existence of a conformal conjugacy defined between some neighborhoods of their Julia sets in proving (i) $\Rightarrow$ (ii). However, since the existence of such a conjugacy implies that they are externally equivalent, they are analytically conjugate near the *filled* Julia sets [DH85].

In this paper, we do not treat with the precise definition of tameness, which is done in terms of conformal measures, so we do not give it here. See [Urb97] for details (including the next theorem). The important fact is the following.

Theorem 2.5. *Every polynomial-like mapping with no recurrent critical points in its Julia set (abbr. NCP) is tame.*

In this paper, we mainly concern with polynomial-like mappings which are restrictions of (some iterate of) global dynamics. In this case, we can prove much stronger conclusion by the following theorem [Ino08].

Theorem 2.6. *Let f_1 and f_2 be two rational or entire maps. Assume they have polynomial-like restrictions $f_i : U'_i \rightarrow U_i$, $i = 1, 2$ which are analytically conjugate. Then there exist rational or entire maps g , φ_1 and φ_2 such that $\varphi_i \circ g = f_i \circ \varphi_i$ and g has a polynomial-like restriction $g : V' \rightarrow V$ analytically conjugate to $f_i : U'_i \rightarrow U_i$.*

Furthermore, if both f_1 and f_2 are rational (resp. polynomials), then g , φ_1 and φ_2 are also rational (resp. polynomials), i.e., f_1 and f_2 are conjugate by an irreducible holomorphic correspondence. In particular, f_1 and f_2 have the same degree.

Theorem 2.2 is an easy consequence of these theorems.

3. ANALYTIC FAMILIES OF POLYNOMIAL-LIKE MAPPINGS

In this section, we briefly review the notion of analytic family of polynomial-like mappings and its straightening map. We also consider families with marked points.

Definition (AFPL). An *analytic family of polynomial-like mappings* (abbr. AFPL) of degree d is a family $\mathbf{f} = (f_\mu : U'_\mu \rightarrow U_\mu)_{\mu \in \Lambda}$ of polynomial-like mappings of degree d parameterized by a complex manifold Λ such that

- (i) $\mathcal{U} = \{(\mu, z); z \in U_\mu\}$ and $\mathcal{U}' = \{(\mu, z); z \in U'_\mu\}$ are homeomorphic over Λ to $\Lambda \times \Delta$, where Δ is the unit disk;
- (ii) the projection from the closure of $\mathcal{U}'$ in $\mathcal{U}$ to Λ is proper;
- (iii) the map $f : \mathcal{U}' \rightarrow \mathcal{U}$, $f(\mu, z) = (\mu, f_\mu(z))$ is holomorphic and proper.

Let $\mathcal{C}(\mathbf{f}) = \{\mu \in \Lambda; K(f_\mu) \text{ is connected}\}$ be the *connectedness locus* of $\mathbf{f}$.

Definition (External markings for AFPL). Let $\mathbf{f} = (f_\mu : U'_\mu \rightarrow U_\mu)_{\mu \in \Lambda}$ be an AFPL. An *external marking* for $\mathbf{f}$ is a family of accesses $([\gamma_\mu])_{\mu \in \mathcal{C}(\mathbf{f})}$ such that $(\mu, t) \mapsto \gamma_\mu(t)$ is continuous for $(\mu, t) \in \mathcal{C}(\mathbf{f}) \times (0, 1]$.

An *externally marked AFPL* is a pair $(\mathbf{f}, [\gamma_\mu])$ of an AFPL and an external marking for it.

Notice that we only consider external markings for maps with connected Julia sets.

Remark 3.1. We do not require that $(\mu, t) \mapsto \gamma_\mu(t)$ is continuous on $\mathcal{C}(\mathbf{f}) \times [0, 1]$. Indeed, consider the quadratic family $\mathbf{Q} = (Q_c(z) = z^2 + c)_{c \in \mathbb{C}}$ with standard external marking. Namely, let $\gamma_c(t) = \phi_c^{-1}(\exp(t))$ where ϕ_c is the Böttcher coordinate for Q_c , and consider $[\gamma_c]$ as an external marking. Then $(c, t) \mapsto \gamma_c(t)$ is not continuous at $(1/4, 0)$ because of the parabolic implosion (discontinuity of the filled Julia set). Thus it is not reasonable to require continuity at $t = 0$.

By the straightening theorem, we can naturally define a map from the connectedness locus $\mathcal{C}(\mathbf{f})$ for an externally marked AFPL to $\mathcal{C}(d)$.

Definition (Straightening maps for AFPL). Let $\mathbf{f} = (f_\mu : U'_\mu \rightarrow U_\mu)_{\mu \in \Lambda}$ be an AFPL and $\Gamma = [\gamma_\mu]_{\mu \in \mathcal{C}(\mathbf{f})}$ be an external marking. The *straightening map* $\chi_{\mathbf{f}, \Gamma} : \mathcal{C}(\mathbf{f}) \rightarrow \mathcal{C}(d)$ is defined as follows: $\chi_{\mathbf{f}, \Gamma}(\mu) = g_\mu$ if $f_\mu : U'_\mu \rightarrow U_\mu$ is hybrid equivalent to g_μ respecting the external markings (the external marking for g_μ is the standard external marking).

In the following, whenever we consider an AFPL, we fix an external marking for each AFPL. Hence we omit Γ for simplicity and write $\chi_{\mathbf{f}}$ instead of $\chi_{\mathbf{f}, \Gamma}$.

The following theorem is proved by Douady and Hubbard [DH85].

Theorem 3.2. *The straightening map for an AFPL of degree two is continuous and can be extended continuously on Λ .*

This theorem depends on the following lemma and quasiconformal rigidity of quadratic polynomials in the boundary of the Mandelbrot set (see Theorem 7.2). Namely, for any $f \in \partial\mathcal{C}(2)$, if $g \in \mathcal{C}(2)$ is quasiconformally conjugate to f , then $g = f$.

Lemma 3.3. *Consider an analytic family $(f_\mu : U'_\mu \rightarrow U_\mu)_{\mu \in \Lambda}$ of polynomial-like mappings of degree d and let $\chi : \mathcal{C}(\Lambda) \rightarrow \mathcal{C}(d)$ be its straightening map.*

Assume $\mu_n \rightarrow \mu$ in $\mathcal{C}(\Lambda)$ and $\chi(\mu_n)$ converges to some $P \in \mathcal{C}(d)$. Then there exist $K \geq 1$ independent of n and a K -quasiconformal hybrid conjugacy ψ_n between f_{μ_n} and $\chi(\mu_n)$ such that ψ_n converges uniformly to a K -quasiconformal conjugacy ψ between f_μ and P by passing to a subsequence. In particular, $\chi(\mu)$ and P are quasiconformally equivalent.

However, quasiconformal rigidity does not hold for polynomials of higher degree in the bifurcation locus. For example, if the basin of a parabolic periodic orbit contains two or more critical points with distinct grand orbits, then you can deform it quasiconformally to another polynomial in $\partial\mathcal{C}(d)$. Discontinuity of straightening maps is caused by such a lack of quasiconformal rigidity and Douady and Hubbard used such a parabolic polynomial to construct an example of discontinuous straightening map [DH85].

Definition (Marked points, AFPL(n)MP). Let $\mathbf{f}_0 = (f_\mu : U'_\mu \rightarrow U_\mu)$ be an AFPL. A *marked point* x_μ for $\mathbf{f}$ is a holomorphic map $x : \Lambda \rightarrow \mathbb{C}$ such that $x_\mu = x(\mu) \in U_\mu$.

An *analytic family of polynomial-like mappings with a marked point* (abbr. AFPLMP) is a family

$$\mathbf{f} = (f_\mu : U'_\mu \rightarrow U_\mu, x_\mu)_{\mu \in \Lambda}$$

such that $\mathbf{f}_0 = (f_\mu : U'_\mu \rightarrow U_\mu)$ is an AFPL and x_μ is a marked point for $\mathbf{f}$.

Let us denote

$$\mathcal{CH}(\mathbf{f}) = \{\mu \in \Lambda; K(f_\mu) \text{ is connected and } x_\mu \in K(f_\mu)\}.$$

The *straightening map* $\chi_{\mathbf{f}} : \mathcal{CH}(\mathbf{f}) \rightarrow \mathcal{CH}(d)$, where

$$\mathcal{CH}(d) = \{(g, z); g \in \mathcal{C}(d) \text{ and } z \in K(g)\} \subset \text{Poly}(d) \times \mathbb{C},$$

is defined as follows. Let $\chi_{\mathbf{f}}(\mu) = (g_{\mu}, z_{\mu})$ when $f_{\mu} : U'_{\mu} \rightarrow U_{\mu}$ is hybrid equivalent to g_{μ} by a hybrid conjugacy ψ and $\psi(x_{\mu}) = z_{\mu}$ (note that $\psi|_{K(f_{\mu})}$ is unique under the assumption that ψ respects external markings).

We need also consider an *analytic family of polynomial-like mappings with two marked points* (abbr. *AFPL2MP*). More generally, for $n \geq 1$, we say a family

$$\mathbf{f} = (f_{\mu} : U'_{\mu} \rightarrow U_{\mu}, x_{1,\mu}, \dots, x_{n,\mu})_{\mu \in \Lambda}$$

is an *AFPLnMP* if $\mathbf{f}_0 = (f_{\mu})_{\mu \in \Lambda}$ is an AFPL and $x_1, \dots, x_n$ are marked points for $\mathbf{f}_0$. (equivalently, $\mathbf{f}_k = (f_{\mu}, x_{k,\mu})_{\mu \in \Lambda}$ is AFPLMP for any $k = 1, \dots, n$). We can similarly define the *straightening map* as follows: Let

$$\mathcal{CH}(\mathbf{f}) = \bigcap_{k=1}^n \mathcal{CH}(\mathbf{f}_k),$$

$$\mathcal{CH}^n(d) = \{(g, z_1, \dots, z_n); g \in \mathcal{C}(d), z_k \in K(g) \text{ for } k = 1, \dots, n\},$$

and define $\chi_{\mathbf{f}} : \mathcal{CH}(\mathbf{f}) \rightarrow \mathcal{CH}^n(d)$ by $\chi_{\mathbf{f}}(\mu) = (g, z_1, \dots, z_n)$ when $\chi_{\mathbf{f}_k}(\mu) = (g, z_k)$.

We will discuss continuity of straightening maps for AFPL2MP in Section 5.1.

4. PARABOLIC IMPLOSION

Here we recall the notion of parabolic implosion and geometric limit. For more details, see [Dou94], [DSZ97], [Shi98] and [Shi00].

Let f_0 be a holomorphic map defined near 0. Assume 0 is a non-degenerate 1-parabolic fixed point, that is, $f_0(0) = 0$, $f'_0(0) = 1$ and $f''_0(0) \neq 0$. By a change of coordinate, we may assume f_0 has the form

$$f_0(z) = z + z^2 + O(z^3)$$

as $z \rightarrow 0$. For $\varepsilon > 0$, consider two disks

$$D_{f_0, \text{attr}} = \{z \in \mathbb{C}; |z + \varepsilon| < \varepsilon\}, \quad D_{f_0, \text{rep}} = \{z \in \mathbb{C}; |z - \varepsilon| < \varepsilon\}.$$

If ε is sufficiently small, then

$$f(D_{f_0, \text{attr}}) \subset D_{f_0, \text{attr}}, \quad f(D_{f_0, \text{rep}}) \supset D_{f_0, \text{attr}},$$

and there exist conformal maps

$$\Phi_{f_0, \text{attr}} : D_{f_0, \text{attr}} \rightarrow \mathbb{C}, \quad \Phi_{f_0, \text{rep}} : D_{f_0, \text{rep}} \rightarrow \mathbb{C}$$

satisfying the Abel equation:

$$(1) \quad \Phi_{f_0, *}(f_0(z)) = \Phi_{f_0, *}(z) + 1 \quad (* = \text{attr}, \text{rep}),$$

where both sides are defined. We call $\Phi_{f_0, \text{attr}}$ (resp. $\Phi_{f_0, \text{rep}}$) an *attracting Fatou coordinate* (resp. *repelling Fatou coordinate*) for f_0 . They are unique up to post-composition by translation. If f is a rational map or an entire map, we can extend Fatou coordinates by the functional equation (1):

- The domain of $\Phi_{f_0, \text{attr}}$ can be extended to the whole basin of attraction B_0 of 0.
- The domain of $\Psi_{f_0, \text{rep}} = \Phi_{f_0, \text{rep}}^{-1}$ can be extended to the whole complex plane $\mathbb{C}$.

For $c \in \mathbb{C}$, let us define $g_{f_0, c} : B_0 \rightarrow \mathbb{C}$ by

$$g_{f_0, c}(z) = \Phi_{f_0, \text{rep}}^{-1}(\Phi_{f_0, \text{attr}}(z) + c).$$

Then $g_{f_0, c}$ commutes with f_0 , i.e., $g_{f_0, c} \circ f_0 = f_0 \circ g_{f_0, c}$. We call $g_{f_0, c}$ a *Lavaurs map* of f and we call c the *phase* of $g_{f_0, c}$.

Let f be a small perturbation of f_0 . By taking an affine conjugacy, we may assume 0 is still a fixed point for f . Let us denote $f'(0) = \exp(2\pi i\alpha)$ with α small. Here we consider the case $\alpha \neq 0$ and $|\arg \alpha| < \pi/4$ (or $|\arg(-\alpha)| < \pi/4$). Let x be the other fixed point for f close to 0 (bifurcated from 0). Then $x = -2\pi i\alpha(1 + o(1))$ as $f \rightarrow f_0$. Let $D_{f,\text{attr}}$ and $D_{f,\text{rep}}$ be the disks of radii ε whose boundaries pass through 0 and x such that $D_{f,\text{attr}}$ intersects the negative real axis and $D_{f,\text{rep}}$ intersects the positive real axis, so that $D_{f,*} \rightarrow D_{f_0,*}$ as $f \rightarrow f_0$. Then there exists a conformal map Φ_f defined on $D_{f,\text{attr}} \cup D_{f,\text{rep}}$ such that $\Phi_f(f(z)) = \Phi_f(z) + 1$. We call Φ_f a *Fatou coordinate* for f . It is also unique up to post-composition by translation. Fatou coordinates depend continuously on f if we normalize properly. More precisely, what we need is the following fact: If $f_n \rightarrow f_0$, then there exist sequences c_n and C_n of complex numbers such that

$$\Phi_{f_n}(z) + c_n \rightarrow \Phi_{f_0,\text{attr}}(z) \text{ on } D_{f_0,\text{attr}}, \quad \Phi_{f_n}(z) + C_n \rightarrow \Phi_{f_0,\text{rep}}(z) \text{ on } D_{f_0,\text{rep}}.$$

as $n \rightarrow \infty$. Hence we have

$$\begin{aligned} f_n^k &= \Phi_{f_n}^{-1}(\Phi_{f_n}(z) + k) \\ &= \Phi_{f_n}^{-1}(\Phi_{f_n}(z) + c_n + (k - c_n + C_n) - C_n). \end{aligned}$$

Now assume $c_n - C_n$ converges in $\mathbb{C}/\mathbb{Z}$, i.e., there exists a sequence $k_n \in \mathbb{Z}$ such that $\lim_{n \rightarrow \infty} k_n - c_n + C_n = c \in \mathbb{C}$. Then we get a convergence

$$f_n^{k_n}(z) \rightarrow g_{f_0,c}(z).$$

We say that f_n converges geometrically to $(f_0, g_{f_0,c})$ and denote

$$f_n \xrightarrow{\text{geom}} (f_0, g_{f_0,c}).$$

5. CONTINUOUS STRAIGHTENING MAPS AND MULTIPLIERS

The following theorem relates continuity of straightening map to a condition on multipliers, and is the key to get discontinuity of straightening maps.

Theorem 5.1. *Let $\mathbf{f} = (f_\mu : U'_\mu \rightarrow U_\mu, x_\mu, y_\mu)_{\mu \in \Lambda}$ be an AFPL2MP of degree $d \geq 2$. Assume*

- (i) *for any $\mu \in \Lambda$, 0 is a fixed point for f_μ ;*
- (ii) *α_μ is a marked repelling periodic point;*
- (iii) *for $\mu = \mu_0$, 0 is a non-degenerate 1-parabolic fixed point, $x_{\mu_0} = y_{\mu_0}$ and they lie in the basin of 0 for f_{μ_0} ;*
- (iv) *there exist sequences $\mu_n \xrightarrow{n \rightarrow \infty} \mu_0$ and $\mu_{n,m} \xrightarrow{m \rightarrow \infty} \mu_n$ such that*
 - $\mu_n, \mu_{n,m} \in \mathcal{C}(\mathbf{f})$;
 - $x_{\mu_n} \neq y_{\mu_n}$ for $n \geq 1$;
 - 0 is a non-degenerate 1-parabolic fixed point for f_{μ_n} ;
 - $f_{\mu_{n,m}} \xrightarrow{\text{geom}} (f_{\mu_n}, g_n)$ as $m \rightarrow \infty$ for some Lavaurs map g_n such that $g_n(x_{\mu_n}) = \alpha_{\mu_n}$. In particular, 0 is no more a parabolic fixed point for $f_{\mu_{n,m}}$.
 - $g_n \rightarrow g$ for some Lavaurs map g for f_{μ_0} such that $g'(x_{\mu_0}) \neq 0$.
- (v) $\chi_{\mathbf{f}}(\mu_{n,m}) \rightarrow \chi_{\mathbf{f}}(\mu_n)$ as $m \rightarrow \infty$ and $\chi_{\mathbf{f}}(\mu_n) \rightarrow \chi_{\mathbf{f}}(\mu_0)$ as $n \rightarrow \infty$.

Then

$$|\text{mult}_{f_{\mu_0}}(\alpha_{\mu_0})| = |\text{mult}_{P_{\mu_0}}(\psi_{\mu_0}(\alpha_{\mu_0}))|,$$

where $\chi_{\mathbf{f}}(\mu) = (P_\mu, x_\mu^P, y_\mu^P)$ and ψ_μ is a hybrid conjugacy between f_μ and P_μ .

Roughly speaking, if the moduli of the multipliers of the corresponding repelling periodic points α_{μ_0} and $\alpha_{\mu_0}^P$ are different and there are plenty of perturbations in $\mathcal{C}(\mathbf{f})$, then the straightening map $\chi_{\mathbf{f}}$ is discontinuous.

The rest of this section is devoted to proof this theorem. We may assume that the hybrid conjugacy $\psi_{\mu_n, m}$ converges a quasiconformal conjugacy φ_n between f_{μ_n} and P_{μ_n} as $m \rightarrow \infty$ by Lemma 3.3. Then by the continuity (v), we have

$$\varphi_n(x_{\mu_n}) = \lim_{m \rightarrow \infty} \psi_{\mu_n, m}(x_{\mu_n, m}) = \psi_{\mu_n}(x_{\mu_n}) = x_{\mu_n}^P,$$

and similarly we have $\varphi_n(y_{\mu_n}) = y_{\mu_n}^P$.

On the other hand, since $f_{\mu_n, m} \xrightarrow{\text{geom}} (f_{\mu_n}, g_n)$, there exists a sequence $(k_{n, m})$ such that

$$f_{\mu_n, m}^{k_{n, m}} \rightarrow g_n.$$

This implies that, if w satisfies that $f_{\mu_n, m}^{k_{n, m}}(\psi_{\mu_n, m}^{-1}(w))$ lies in the definition of φ_n for sufficiently large n , we have

$$P_{\mu_n, m}^{k_{n, m}}(w) = \psi_{\mu_n, m} \circ f_{\mu_n, m}^{k_{n, m}} \circ \psi_{\mu_n, m}^{-1}(w) \rightarrow g_n^P(w) = \varphi_n \circ g_n \circ \varphi_n^{-1}(w),$$

namely, $P_{\mu_n, m}$ geometrically converges to (P_{μ_n}, g_n^P) . Then

$$\begin{aligned} g_n^P \circ \psi_{\mu_n}(x_{\mu_n}) &= g_n^P \circ \varphi_n(x_{\mu_n}) \\ &= \varphi_n(g_n(x_{\mu_n})) \\ &= \varphi_n(\alpha_{\mu_n}) \end{aligned}$$

is a repelling periodic point for P_{μ_n} . Let us denote it by $\alpha_{\mu_n}^P$. Then $\alpha_{\mu_n}^P = \psi_{\mu_n}(\alpha_{\mu_n})$. In fact, the proof of [DH85, p. 302, Lemma 1] can be applied to our case to show that $\psi_{\mu_n} = \varphi_n$ on the Julia set. Observe that the combinatorial assumption there holds because the images of the unique parabolic basin of 0 by them are the same.

Now let

$$\delta_n(w) = \log \left| \frac{\varphi_n(w) - \alpha_{\mu_n}^P}{w - \alpha_{\mu_n}} \right|.$$

By passing to a further subsequence, we may assume φ_n also converges as $n \rightarrow \infty$. Then we have the following “distortion” property for φ_n at α_{μ_n} :

Lemma 5.2. *Let $a_n = |\text{mult}_{f_{\mu_n}}(\alpha_{\mu_n})|$ and $b_n = |\text{mult}_{P_{\mu_n}}(\alpha_{\mu_n}^P)|$. Then*

$$(2) \quad \delta_n(w) = \frac{\log b_n - \log a_n}{\log a_n} \log |w - \alpha_{\mu_n}| + O(1)$$

as $w \rightarrow \alpha_{\mu_n}$ uniformly on n . In particular, if $|\text{mult}_{f_{\mu_0}}(\alpha_{\mu_0})| \neq |\text{mult}_{P_{\mu_0}}(\alpha_{\mu_0}^P)|$, then $\delta_n(w)$ diverges as $w \rightarrow \alpha_{\mu_n}$ uniformly on sufficiently large n .

Remark 5.3. This Lemma is equivalent that the Hölder exponent of φ_n at α_{μ_n} is equal to $\log b_n / \log a_n$, i.e.,

$$|\varphi_n(w) - \varphi_n(\alpha_{\mu_n})| \asymp |w - \alpha_{\mu_n}|^{\frac{\log b_n}{\log a_n}}.$$

Proof. Take a small circle $S(r, \alpha_{\mu_n})$ centered at α_{μ_n} . If the radius $r > 0$ is sufficiently small, then the circle and its image $f_{\mu_n}^P(S(r, \alpha_{\mu_n}))$ bounds an annulus A , where p is the period of

α_{μ_n} for f_{μ_n} . There exists some constant $C > 1$ such that $C^{-1} < |\delta_n(w)| < C$ for any $w \in \bar{A}$. For w close to α_{μ_n} , there exists some $k > 0$ such that $f_{\mu_n}^{kp}(w) \in \bar{A}$. Then

$$\begin{aligned} \delta_n(w) &= \log \left| \frac{\varphi_n(w) - \varphi_n(\alpha_{\mu_n})}{P_{\mu_n}^{kp}(\varphi_n(w)) - \varphi_n(\alpha_{\mu_n})} \right| + \log \left| \frac{\varphi_n(f_{\mu_n}^{kp}(w)) - \varphi_n(\alpha_{\mu_n})}{f_{\mu_n}^{kp}(w) - \alpha_{\mu_n}} \right| + \log \left| \frac{f_{\mu_n}^{kp}(w) - \alpha_{\mu_n}}{w - \alpha_{\mu_n}} \right| \\ &= \eta_1(w) + \delta_n(f_{\mu_n}^{kp}(w)) + \eta_2(w). \end{aligned}$$

Since $f_{\mu_n}^p$ and $P_{\mu_n}^p$ are linearizable near α_{μ_n} and $\alpha_{\mu_n}^p = \varphi_n(\alpha_{\mu_n})$ respectively, it follows that $\eta_1(w) = -k \log a_n + O(1)$ and $\eta_2(w) = k \log b_n + O(1)$. Therefore, $\delta_n(w) = k(\log b_n - \log a_n) + O(1)$. Furthermore, as $w \rightarrow \alpha_{\mu_n}$, k tends to infinity, so we have (2). These estimates are uniform on n because of the convergence as $n \rightarrow \infty$. $\square$

Let $w = g_n(y_{\mu_n})$. If n is sufficiently large, w is close to α_{μ_n} , thus we have

$$\begin{aligned} \delta_n(w) &= \log \left| \frac{\varphi_n(g_n(y_{\mu_n})) - \varphi_n(g_n(x_{\mu_n}))}{g_n(y_{\mu_n}) - g_n(x_{\mu_n})} \right| \\ &= \log \left| \frac{g_n^p \circ \varphi_n(y_{\mu_n}) - g_n^p \circ \varphi_n(x_{\mu_n})}{y_{\mu_n} - x_{\mu_n}} \right| - \log \left| \frac{g_n(y_{\mu_n}) - g_n(x_{\mu_n})}{y_{\mu_n} - x_{\mu_n}} \right| \\ &= \log \left| \frac{g_n^p \circ \psi_n(y_{\mu_n}) - g_n^p \circ \psi_n(x_{\mu_n})}{y_{\mu_n} - x_{\mu_n}} \right| - \log \left| \frac{g_n(y_{\mu_n}) - g_n(x_{\mu_n})}{y_{\mu_n} - x_{\mu_n}} \right|. \end{aligned}$$

Since x_{μ_n} and y_{μ_n} lie in the interior of the filled Julia set, where ψ_n is holomorphic, we have

$$\delta_n(w) = \log \left(\frac{(g_n^p \circ \psi_n)'(x_{\mu_n})}{g_n'(x_{\mu_n})} + O(|y_n - x_n|) \right).$$

Since $g_n \rightarrow g$ by assumption, we may assume that $g_n^p = \varphi_n \circ g_n \circ \varphi_n^{-1}$ also converges by passing to a subsequence. This implies that $|\delta_n(w)|$ is bounded uniformly for sufficiently large n .

Therefore, by Lemma 5.2, this holds only when $|\text{mult}_{f_{\mu_0}}(\alpha_{\mu_0})| = |\text{mult}_{P_{\mu_0}}(\psi_{\mu_0}(\alpha_{\mu_0}))|$. This proves the theorem. $\square$

6. COMBINATORICS OF DYNAMICS OF POLYNOMIALS

We need to find nice perturbations of a given parabolic polynomials to apply Theorem 5.1 to a family of renormalizable polynomials. To do this end, we need most of the results in the preceding paper [IK08] with Kiwi. One of the most essential tools to construct such perturbations is a combinatorial technique which we call *combinatorial tuning*, but we also need injectivity of straightening maps, compactness of renormalizable sets for primitive combinatorics, and so on. Hence we recall some definitions and results in [IK08] in this section. We also prove some lemmas for later use.

6.1. Mapping schemata and skew products. The notion of mapping schema is introduced by Milnor [MP92] to describe the dynamics of hyperbolic polynomials. Here, we review the notion of mapping schemata and consider polynomials and polynomial-like mappings over them, which are simple generalization of usual polynomials and polynomial-like mappings.

Definition (Mapping schemata). A *mapping schema* is a triple $T = (|T|, \sigma, \delta)$ where $|T|$ is a finite set, and $\sigma : |T| \rightarrow |T|$ and $\delta : |T| \rightarrow \mathbb{N}$ are maps such that for any periodic point

$v \in |T|$ for σ , we have

$$\prod_{k=0}^{n-1} \delta(\sigma^k(v)) \geq 2,$$

where n is the period of v . We call δ the *degree function* of T and

$$\delta(T) = 1 + \sum_{v \in |T|} (\delta(v) - 1)$$

the *total degree* of T .

We call $v \in |T|$ is *critical* if $\delta(v) > 1$. We say T is *reduced* if all $v \in |T|$ are critical. Here we only consider reduced mapping schemata because we can easily extract a reduced schema from a given schema by taking the first return map [MP92].

Definition. We say T has a *non-trivial critical relation* if either

- there exist critical $v, v' \in |T|$ and $n > 0$ such that $v \neq v'$ and $v = \sigma^n(v')$, or
- there exists a critical $v \in |T|$ such that $\delta(v) \geq 3$.

Otherwise, we call T is of *disjoint type*. It is equivalent that there is exactly $\delta(T) - 1$ periodic orbits, or every critical $v \in |T|$ is periodic and satisfy

$$\prod_{n=0}^{p-1} \delta(\sigma^n(v)) = 2$$

where p is the period of v .

An integer $d (\geq 2)$ represents the trivial schema of total degree d , i.e., $d = (\{pt\}, id, d)$ (see Figure 1). Another important example of mapping schemata is the schema of capture type: Let $T_{\text{cap},d} = (|T_{\text{cap}}|, \sigma_{\text{cap}}, \delta_{\text{cap},d})$ be defined by

- $|T_{\text{cap}}| = \{v_1, v_2\}$,
- $\sigma_{\text{cap}}(v_j) = v_1$ for $j = 1, 2$,
- $\delta_{\text{cap},d}(v_1) = 2$ and $\delta_{\text{cap},d}(v_2) = d - 1$ for $j = 1, 2$.

For simplicity, we denote by T_{cap} the degree 3 capture schema $T_{\text{cap},3}$.



FIGURE 1. The trivial schema (left) and the capture schema $T_{\text{cap},d}$ (right) of degree d .

Definition (Polynomials over mapping schemata and universal polynomial model spaces). Let $T = (|T|, \sigma, \delta)$ be a mapping schema. A *polynomial over T* is a map $f : |T| \times \mathbb{C} \rightarrow |T| \times \mathbb{C}$ of the form $f(v, z) = (\sigma(v), f_v(z))$ such that f_v is a polynomial of degree $\delta(v)$. We say f is *monic centered* if f_v is so for all $v \in |T|$. The *universal polynomial model space* $\text{Poly}(T)$ is the set of all monic centered polynomials over T .

For a polynomial f over T , the *filled Julia set* $K(f)$ is the set of points whose forward orbit is precompact, and the *Julia set* $J(f)$ is the boundary of $K(f)$. We say $K(f)$ is *fiberwise connected* if the fiber $K(f, v) = \{z \in \mathbb{C}; (v, z) \in K(f)\}$ is connected for all $v \in |T|$. The *(fiberwise) connectedness locus* $\mathcal{C}(T)$ is the set of all maps $f \in \text{Poly}(T)$ with fiberwise connected filled Julia set.

Observe that a polynomial over the trivial schema d is simply a polynomial of degree d . Thus the definition of $\text{Poly}(d)$ is consistent and we can treat normal polynomials and polynomials over mapping schemata at the same time.

For $f \in \text{Poly}(T)$, $v \in |T|$ and $n > 0$, define f_v^n by the equation

$$f^n(v, z) = (\sigma^n(v), f_v^n(z)).$$

Then we have

$$K(f) = \{(v, z); \{f_v^n(z)\}_{n \geq 0} \text{ is bounded}\}.$$

For a polynomial $f \in \text{Poly}(T)$ over a mapping schema $T = (|T|, \sigma, \delta)$, there exists the Böttcher coordinate ϕ_f at $|T| \times \{\infty\}$, i.e., ϕ_f is a holomorphic map defined on a neighborhood of $|T| \times \{\infty\}$ such that

- ϕ_f is tangent to the identity at $|T| \times \{\infty\}$;
- it has the form $\phi_f(v, z) = (v, \phi_{f,v}(z))$;
- it conjugates f to $(v, z) \mapsto (\sigma(v), z^{\delta(v)})$.

If $f \in \mathcal{C}(T)$, then we can extend ϕ_f using the dynamics and obtain a univalent map

$$\phi_f : (|T| \times \mathbb{C}) \setminus K(f) \rightarrow |T| \times (\mathbb{C} \setminus \bar{\Delta}),$$

which we still denote by ϕ_f . Hence for $v \in |T|$ and $\theta \in \mathbb{R}/\mathbb{Z}$, we can define the *external ray* by

$$R_f(v, \theta) = \phi_f^{-1}\{(v, r \exp(2\pi i \theta)); r > 1\}.$$

By definition, we have $f(R_f(v, \theta)) = R_f(\sigma(v), \delta(v)\theta)$. Many results on external rays for usual polynomial also hold for polynomials over mapping schema and proofs are straightforward. For example, every external ray of a rational angle lands at a repelling or parabolic eventually periodic point.

Definition (Polynomial-like mappings over mapping schemata). Let $T = (|T|, \sigma, \delta)$ be a mapping schema. A *polynomial-like mapping over T* is a proper holomorphic skew product over σ

$$g : \begin{array}{ccc} U' & \rightarrow & U \\ (v, z) & \mapsto & (\sigma(v), g_v(z)), \end{array}$$

such that

- $U' \subseteq U$ are subsets of $|S| \times \mathbb{C}$ having the form

$$U' = \bigcup_{v \in |T|} \{v\} \times U'_v, \quad U = \bigcup_{v \in |T|} \{v\} \times U_v,$$

where U'_v and U_v are topological disks;

- g has the form $g(v, z) = (\sigma(v), g_v(z))$ where the degree of $g_v : U'_v \rightarrow U_{\sigma(v)}$ is equal to $\delta(v)$.

We may also write it as a collection of proper holomorphic maps

$$g = (g_v : U'_v \rightarrow U_{\sigma(v)})$$

between topological disks in the complex plane.

The *filled Julia set* $K(g)$ is defined as follows:

$$K(g) = \bigcap_{n \geq 0} g^{-n}(U').$$

and the Julia set $J(g)$ is the boundary of $K(g)$. We say $K(g)$ is *fiberwise connected* if $K(g) \cap \{v\} \times \mathbb{C}$ is connected for all $v \in |T|$. We denote

$$K(g, v) = \{z \in \mathbb{C}; (v, z) \in K(g)\}.$$

By definition, $K(g)$ is fiberwise connected if and only if $K(g, v)$ is connected for all $v \in |T|$.

Definition (External markings). An *external marking* of a polynomial-like mapping g over a mapping schema $T = (|T|, \sigma, \delta)$ is a collection of accesses $([\gamma_v])_{v \in |T|}$ such that $\gamma_v \subset U'_v$ and $f(\gamma_v) \cap U'_{\sigma(v)} \in [\gamma_{\sigma(v)}]$. An *externally marked polynomial-like mapping* over T is a pair $(g, ([\gamma_v]))$ of a polynomial-like mapping over T and an external marking of it.

Let $f \in \mathcal{C}(T)$. The *standard external marking* of f is the external marking $([R_f(v, 0)])_{v \in |T|}$, defined by the external rays of angle zero.

Definition (Hybrid equivalence). Two polynomial-like mappings g_1 and g_2 over a mapping schema $T = (|T|, \sigma, \delta)$ are *hybrid equivalent* if there exists a quasiconformal map ψ defined on a neighborhood of $K(g_1)$ such that $\psi \circ g_1 = g_2 \circ \psi$ where both sides are defined and $\frac{\partial \psi}{\partial z} \equiv 0$ a.e. on $K(g_0)$.

When g_1 and g_2 are externally marked, we say that a hybrid conjugacy ψ *preserves external markings* if the external marking of g_1 is mapped to that of g_2 by ψ .

We can generalize the straightening theorem (Theorem 2.1) to this case [IK08, Theorem A].

Theorem 6.1 (Straightening theorem for polynomial-like mappings over mapping schemata). *A polynomial-like mapping g over a mapping schema $T = (|T|, \sigma, \delta)$ is hybrid equivalent to some $f \in \text{Poly}(T)$. Furthermore, if $K(g)$ is fiberwise connected and g is externally marked, then there exists a unique $f \in \mathcal{C}(d)$ such that a hybrid conjugacy between f and g preserves external markings, where the external marking of f is the standard external marking.*

6.2. Rational laminations. For an integer $d > 1$, let m_d denote the d -fold covering $\theta \mapsto d\theta$ defined on $\mathbb{R}/\mathbb{Z}$ to itself. For a mapping schema $T = (|T|, \sigma, \delta)$, define a map $m_T : |T| \times \mathbb{R}/\mathbb{Z} \rightarrow |T| \times \mathbb{R}/\mathbb{Z}$ by

$$m_T(v, \theta) = (\sigma(v), m_{\delta(v)}(\theta)) = (\sigma(v), \delta(v)\theta).$$

We say two sets $A, B \subset \mathbb{R}/\mathbb{Z}$ (or $\{v\} \times \mathbb{R}/\mathbb{Z}$) are *unlinked* if B is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus A$. Note that it is equivalent that A is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus B$, or that the Euclidean (or hyperbolic) convex hulls of $\exp(2\pi iA)$ and $\exp(2\pi iB)$ in $\mathbb{C}$ are disjoint.

Let $A \subset \mathbb{R}/\mathbb{Z}$. We say a map $f : A \rightarrow f(A) \subset \mathbb{R}/\mathbb{Z}$ is *consecutive preserving* if for any component (θ, θ') of $\mathbb{R}/\mathbb{Z} \setminus A$, $(f(\theta), f(\theta'))$ is a component of $\mathbb{R}/\mathbb{Z} \setminus f(A)$.

Definition (Invariant rational laminations). Let $T = (|T|, \sigma, \delta)$ be a mapping schema. An equivalence relation λ on $|T| \times \mathbb{Q}/\mathbb{Z}$ is called a *T -invariant rational lamination* or a *rational lamination over T* if the following conditions hold:

- (i) Each equivalence class is contained in $\{v\} \times \mathbb{Q}/\mathbb{Z}$ for some $v \in |T|$.
- (ii) λ is closed in $(|T| \times \mathbb{Q}/\mathbb{Z})^2$.
- (iii) Every equivalence class is finite.
- (iv) Equivalence classes are pairwise unlinked.
- (v) For a λ -equivalence class A , $m_T(A)$ is also a λ -equivalence class.
- (vi) $m_T : A \rightarrow m_T(A)$ is consecutive preserving.

Let us denote by $\text{supp}(\lambda) \subset |T| \times \mathbb{Q}/\mathbb{Z}$ the union of all non-trivial λ -classes.

We may denote λ as a collection $(\lambda_v)_{v \in |T|}$ of (non-invariant) rational laminations on $\mathbb{Q}/\mathbb{Z}$, i.e., (v, θ) and (v, θ') are λ -equivalent if and only if θ and θ' are λ_v -equivalent.

Example. For $f \in \mathcal{C}(T)$, the *rational lamination* λ_f of f is the landing relation of external rays of rational angles. Namely, (v, θ) and (v, θ') are λ_f -equivalent if and only if the external rays $R_f(v, \theta)$ and $R_f(v, \theta')$ land at the same point. By the theorem of Kiwi [Kiw01], an equivalence relation λ on $|T| \times \mathbb{Q}/\mathbb{Z}$ is a T -invariant rational lamination if and only if λ is the rational lamination of some $f \in \mathcal{C}(T)$.

Definition (Combinatorial renormalization). We say a T -invariant rational lamination λ is *admissible* for $f \in \mathcal{C}(T)$ if $\lambda \subset \lambda_f$. We also say that f is λ -combinatorially renormalizable or f *admits* λ . Namely, $f \in \mathcal{C}(T)$ if and only if $R_f(v, \theta)$ and $R_f(v, \theta')$ land at the same point when (v, θ) and (v, θ') are λ -equivalent. Let

$$\mathcal{C}(\lambda) = \{f \in \mathcal{C}(T); \lambda \subset \lambda_f\}$$

be the set of all polynomials which admit λ .

A rational lamination λ naturally induces the *unlinked relation* for irrational angles, which is closely related to *gaps* introduced by Thurston [Thu85].

Definition (Unlinked classes). Let T be a mapping schema and let λ be a T -invariant rational lamination. We say $(v, \theta), (v', \theta') \in |T| \times (\mathbb{R} \setminus \mathbb{Q})/\mathbb{Z}$ are λ -unlinked if $v = v'$ and for any λ -equivalence class $\{v\} \times A$, θ and θ' lie in the same component of $\mathbb{R}/\mathbb{Z} \setminus A$.

Observe that λ -unlinked relation is an equivalence relation and each equivalence class (λ -unlinked class) is contained in $v \times (\mathbb{R} \setminus \mathbb{Q})/\mathbb{Z}$ for some $v \in |T|$. A set $\{v\} \times L$ is an unlinked class if and only if L is a λ_v -unlinked class.

Lemma 6.2. *Let λ_0 and λ be rational laminations and assume $\lambda \supset \lambda_0$. If a λ -equivalence class A is not a λ_0 -class, then there exists some λ_0 -unlinked class L such that A and L are linked (not unlinked) and $\partial L \cap A$ is nonempty.*

Proof. Let $B \subset A$ be a λ_0 -class. Take a component (s, t) of $\mathbb{R}/\mathbb{Z} \setminus B$ which intersects A . Consider a set

$$F = [s, t] \setminus \bigcup_{\theta, \theta'} [\theta, \theta'],$$

where the union is taken for all λ_0 -equivalent pairs θ, θ' such that $[\theta, \theta'] \subset (s, t)$. Since each pair of such intervals are either disjoint or one contains the other, F is a Cantor set removing countably many points. In particular, F is uncountable and contained in the derived set of F itself. Hence $L = F \cap (\mathbb{R} \setminus \mathbb{Q})/\mathbb{Z}$ is nonempty. Furthermore, since L is unlinked with any λ_0 -equivalence class, L is in fact a λ_0 -unlinked class.

Let $B' \subset A \cap (s, t)$ be another λ_0 -class. Then since B and B' lie in different components of L , A and L are linked.

By construction, $s, t \in A$ lie in the closure of L . □

The external rays for f of λ -equivalent angles cut the phase space into sectors. This allows us to associate a continuum for each λ -unlinked class (compare [Sch04]).

Definition (Sectors and fibers). Let λ be a rational lamination over a mapping schema T and let $f \in \mathcal{C}(\lambda)$. For a λ -unlinked class $L \subset \{v\} \times \mathbb{R}/\mathbb{Z}$ and λ -equivalent angles $(v, \theta), (v, \theta')$, let

$$\text{Sector}_f(v, \theta, \theta'; L)$$

be the connected component of

$$(\{v\} \times \mathbb{C}) \setminus \overline{(R_f(v, \theta) \cup R_f(v, \theta'))}$$

containing the external ray $R_f(v, t)$ for every $(v, t) \in L$. The *fiber* of L is defined by:

$$K_f(L) = K(f) \cap \bigcap_{\theta \sim_{\lambda} \theta', \theta \neq \theta'} \overline{\text{Sector}(v, \theta, \theta'; L)}.$$

The following proposition (see [IK08]) describes some basic properties for λ -unlinked classes and corresponding fibers.

Proposition 6.3. *Let T be a mapping schema and let λ be a T -invariant rational lamination. For $f \in \mathcal{C}(\lambda)$ and a λ -unlinked class L , we have the following:*

- (i) $m_T(L)$ is also a λ -unlinked class.
- (ii) $f(K_f(L)) = K_f(L)$.
- (iii) If L is finite, then
 - (a) $m_T : L \rightarrow m_T(L)$ is a $\delta(L)$ -to-one consecutive preserving map for some $\delta(L) > 0$.
 - (b) $f : K_f(L) \rightarrow K_f(m_T(L))$ has degree $\delta(L)$.
- (iv) If L is infinite, then
 - (a) L is eventually periodic by m_T .
 - (b) There exists a homeomorphism $\alpha_L : \bar{L}/\lambda \rightarrow \mathbb{R}/\mathbb{Z}$ such that $\alpha_{m_T(L)} \circ m_T \circ \alpha_L^{-1}$ is well-defined and coincides with $m_{\delta(L)}$ for some $\delta(L) \geq 1$.

Remark 6.4. We stated this proposition only for the case of d -invariant rational laminations in [IK08], but the same proof can be applied also to rational laminations over mapping schemata.

Similarly, some theorems below in this section is generalized to rational laminations and polynomials over mapping schemata, but the proofs are exactly the same (only notations will be more complicated), and some of them are immediate consequence from the same result for the usual case.

This proposition allows us to describe the dynamics of fibers in terms of λ -equivalence classes and unlinked classes.

Definition (Critical elements). Let λ be a T -invariant rational lamination. For a λ -class A , let $\delta(A)$ denotes the degree of $m_T : A \rightarrow m_T(A)$. It is well-defined by the consecutive preservingness. We say A is *critical* if $\delta(A) > 1$. Similarly, for λ -unlinked class L , $\delta(L)$ is the one defined in Proposition 6.3 we say L is *critical* if $\delta(L) > 1$.

Let $\text{Crit}^P(\lambda)$, $\text{Crit}^W(\lambda)$ and $\text{Crit}^F(\lambda)$ be the set of all critical λ -classes, finite λ -unlinked classes and infinite λ -unlinked classes respectively, and let $\text{Crit}(\lambda) = \text{Crit}^F(\lambda) \cup \text{Crit}^W(\lambda) \cup \text{Crit}^P(\lambda)$. We call an element in $\text{Crit}^P(\lambda) \cup \text{Crit}^W(\lambda)$ (resp. $\text{Crit}^F(\lambda)$) a *Julia critical element*, (resp. a *Fatou critical element*). For a Julia critical element A , we say A is *preperiodic* if $A \in \text{Crit}^P(\lambda)$ and A is *wandering* if $A \in \text{Crit}^W(\lambda)$.

Roughly speaking, critical elements correspond to critical points for $f \in \mathcal{C}(\lambda)$. It follows that

$$\delta(v) - 1 = \sum_{A \in \text{Crit}(\lambda), A \subset (v \times \mathbb{R}/\mathbb{Z})} (\delta(A) - 1).$$

For $* = P, W, F$, let

$$\begin{aligned} PC^*(\lambda) &= \{m_T^n(w); w \in \text{Crit}^*(\lambda), n > 0\}, & PC(\lambda) &= PC^P(\lambda) \cup PC^W(\lambda) \cup PC^F(\lambda), \\ CO^*(\lambda) &= PC^*(\lambda) \cap CO^*(\lambda), & CO(\lambda) &= CO^P(\lambda) \cup CO^W(\lambda) \cup CO^F(\lambda). \end{aligned}$$

Definition (Post-critically finite, hyperbolic and Misiurewicz laminations). We say a T -invariant rational lamination λ is

- *post-critically finite* if there is no wandering critical Julia element (i.e., $\text{Crit}^W(\lambda) = \emptyset$),
- *hyperbolic* if there is no Julia critical elements (i.e., $\text{Crit}(\lambda) = \text{Crit}^F(\lambda)$), and
- *Misiurewicz* if there is no critical λ -unlinked class (i.e., $\text{Crit}(\lambda) = \text{Crit}^P(\lambda)$).

Observe that λ is post-critically finite if and only if $PC(\lambda)$ is finite.

Definition. A T_0 -invariant rational lamination λ is *primitive* if for any infinite λ -unlinked class $w, w' \subset \{v\} \times \mathbb{R}/\mathbb{Z}$, there are no λ -class A such that both $A \cap \overline{w}$ and $A \cap \overline{w'}$ are nonempty.

Even if λ is not primitive, such intersections exist essentially only finitely many.

Lemma 6.5. *Let λ be a post-critically finite rational lamination over a mapping schema T .*

Let A be a λ -class such that there exist infinite λ -unlinked classes $L_1 \neq L_2$ whose closures intersect A . Then there exists some $n \geq 0$ such that either

- $m_T^n(A) \in \text{Crit}^P(\lambda)$, or
- $m_T^n(L_1) \neq m_T^n(L_2)$ and both lie in $CO^F(\lambda)$.

In particular, the set of all eventual periods of such λ -classes A is finite.

Proof. If $m_T^n(A)$ is not critical for any $n \geq 0$, then $m_T^n(L_1) \neq m_T^n(L_2)$ for any $n \geq 0$. Hence the lemma follows because all infinite λ -unlinked classes are eventually periodic. $\square$

Definition (Mapping schemata of rational laminations). Let λ be a T_0 -invariant rational lamination such that $\text{Crit}^F(\lambda)$ is nonempty. Define a (reduced) mapping schema $T(\lambda) = (|T(\lambda)|, \sigma_\lambda, \delta_\lambda)$ by

$$|T(\lambda)| = \text{Crit}^F(\lambda), \quad \sigma_\lambda(w) = m_T^{\ell_w}(w),$$

and $\delta_\lambda = \delta$ is defined in Proposition 6.3, where $\ell_w > 0$ is the smallest number such that $m_T^{\ell_w}(w) \in \text{Crit}^F(\lambda)$.

We say λ has a *non-trivial Fatou critical relation* if $T(\lambda)$ has a non-trivial critical relation. Otherwise, we say λ is of *disjoint type* (it is equivalent that $T(\lambda)$ is of disjoint type).

Proposition 6.3 guarantees the existence of an “internal angle system”, which is needed to let straightening maps well-defined:

Definition (Internal angle systems). Let $T_0 = (|T_0|, \sigma_0, \delta_0)$ be a mapping schema. An *internal angle system* of a T_0 -invariant rational lamination λ is a collection of maps $\alpha = (\alpha_w : \overline{w} \rightarrow \mathbb{R}/\mathbb{Z})_{w \in |T(\lambda)|}$ such that

$$\alpha_{\sigma(w)} \circ m_{T_0}^{\ell_w}(v, \theta) = m_{\delta(w)}(\alpha_{\sigma(w)}(v, \theta)),$$

for $(v, \theta) \in \overline{w}$.

We sometimes omit v and simply write $\alpha_w(\theta)$, for α_w is defined on $\overline{w} \subset \{v\} \times \mathbb{R}/\mathbb{Z}$, so v depends only on w . The map α above can also be considered as follows:

$$\alpha : \bigsqcup_{w \in |T(\lambda)|} \overline{w} \rightarrow |T(\lambda)| \times \mathbb{R}/\mathbb{Z}, \quad \overline{w} \ni (v, \theta) \mapsto \alpha(v, \theta) = (w, \alpha_w(v, \theta)),$$

in this expression, we have

$$m_{T(\lambda)} \circ \alpha(v, \theta) = \alpha \circ m_{T_0}^{\ell_w}(v, \theta)$$

for $(v, \theta) \in \bar{w}$. However, readers should notice that we need to take the disjoint union even if $\bar{w}$ and $\bar{w}'$ intersect for $w \neq w' \in |T(\lambda)|$.

Lemma 6.6. *Let λ be a rational lamination over a mapping schema $T_0 = (|T_0|, \sigma_0, \delta_0)$ and $\alpha = (\alpha_w)_{w \in |T(\lambda)|}$ be an internal angle system. Let $(v, \theta) \in \bar{w}$ be periodic of period p by m_{T_0} . Then $\alpha(v, \theta) = (w, \alpha_w(v, \theta))$ is also periodic of period p' by $m_{T(\lambda)}$, where p' is defined by*

$$\sum_{n=0}^{p'-1} \ell_{\sigma_\lambda^n(w)} = p.$$

In particular, p is not less than the period of w by σ .

Proof. There exists a sequence θ_n such that $(v, \theta_n) \in w$ and either $\theta_n \nearrow \theta$ or $\theta_n \searrow \theta$. It suffices to show when $\theta_n \nearrow \theta$. Then $m_{T_0}^p(v, \theta_n) \nearrow m_{T_0}^p(v, \theta) = (v, \theta)$ and they lie in a λ -unlinked class $w' = \sigma_0^p(w)$. Hence it follows that $w' = w$ and we have

$$\begin{aligned} m_{T(\lambda)}^{p'}(w, \alpha_w(v, \theta)) &= \lim m_{T(\lambda)}^{p'}(w, \alpha_w(v, \theta_n)) \\ &= \lim(\sigma_\lambda^{p'}(w), m_{\delta(\sigma_\lambda^{p'-1}(w))} \circ \cdots \circ m_{\delta(w)} \circ \alpha_w(v, \theta)) \\ &= \lim(\sigma_0^p(w), \alpha_{\sigma_0^p(w)} \circ m_{T_0}^p(v, \theta)) \\ &= (w, \alpha_w(v, \theta)). \end{aligned}$$

On the other hand, let p'_1 be the period of $(w, \alpha_w(v, \theta))$ by $m_{T(\lambda)}$. Then $m_{T_0}^{p'_1}(v, \theta) \sim (v, \theta)$. Moreover,

$$\lim m_{T(\lambda)}^{p'_1}(w, \alpha_w(v, \theta_n)) = m_{T(\lambda)}^{p'_1}(w, \alpha_w(v, \theta)) = (w, \alpha_w(v, \theta)),$$

thus $m_{T_0}^{p'_1}(v, \theta_n) \in \sigma_\lambda^{p'_1}(w) = w$, where $p_1 = \sum_{n=0}^{p'_1-1} \ell_{\sigma_\lambda^n(w)}$. Therefore, w accumulates to $m_{T_0}^{p'_1}(v, \theta)$ from the left, thus $m_{T_0}^{p'_1}(v, \theta) = (v, \theta)$. $\square$

6.3. Renormalizations.

Definition (Renormalizations). Let T_0 be a mapping schema and λ_0 be a T_0 -invariant rational lamination. We say $f \in \mathcal{C}(\lambda_0)$ is λ_0 -renormalizable if there exist topological disks $U'_w \Subset U_w$ for each $w \in |T(\lambda_0)|$ such that

- $g = (f^{\ell_w} : U'_w \rightarrow U_{\sigma(w)})$ is a polynomial-like map over $T(\lambda_0)$ with fiberwise connected Julia set.
- $K(g, w) = K_f(w)$ for all $w \in |T|$.

We call g a λ_0 -renormalization of f .

Definition (Straightening map χ_{λ_0}). Let T_0 be a mapping schema and λ_0 be a T_0 -invariant rational lamination. Let $(\alpha_w : \bar{w} \rightarrow \mathbb{R}/\mathbb{Z})_{w \in |T(\lambda_0)|}$ be an internal angle system. For each $w \in |T(\lambda_0)|$, take $\theta_w \in \bar{w}$ such that $\alpha_w(\theta_w) = 0$.

For $f \in \mathcal{R}(\lambda_0)$, define $\chi_{\lambda_0}(f) \in \text{Poly}(T(\lambda_0))$ as follows; let g be a λ_0 -renormalization of f . Then external rays $([R_f(w, \theta_w)])_{w \in |T(\lambda_0)|}$ defines an external marking of g . Let $\chi_{\lambda_0}(f)$ be the polynomial over $T(\lambda_0)$ hybrid equivalent to g preserving external markings.

This gives a well-defined map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \rightarrow \mathcal{C}(T(\lambda_0))$.

One of the main results in [IK08] is the following:

Theorem 6.7 (Injectivity of straightening maps). *Let T_0 be a mapping schema. For a post-critically finite T_0 -invariant rational lamination λ_0 , the straightening map χ_{λ_0} is injective.*

We can also give some equivalent conditions about the domain $\mathcal{R}(\lambda_0)$ of the straightening map.

Proposition 6.8. *Let λ_0 be a d -invariant rational lamination having nonempty mapping schema $T(\lambda_0)$. Then the following are equivalent:*

- (i) $\mathcal{R}(\lambda_0)$ is nonempty.
- (ii) If A is a critical λ_0 -class and $L \in CO^F(\lambda_0)$, then $A \cap \bar{L} = \emptyset$.

Theorem 6.9 (Compactness of the renormalizable set for primitive combinatorics). *Assume a T_0 -invariant rational lamination λ_0 has nonempty mapping schema $T(\lambda_0)$. Then the following are equivalent:*

- (i) λ_0 is primitive;
- (ii) $\mathcal{C}(\lambda_0) = \mathcal{R}(\lambda_0)$ and it is nonempty;
- (iii) $\mathcal{R}(\lambda_0)$ is compact and nonempty.

To prove a given $f \in \mathcal{C}(\lambda_0)$ is λ_0 -renormalizable, we need the following lemma [IK08, Lemma 6.9], which is based on the idea of “thickening puzzles” by Milnor [Mil00].

Lemma 6.10. *Let λ_0 be a rational lamination over a mapping schema T_0 . Then there exists a proper algebraic set $X \subset \text{Poly}(T_0)$ such that $\mathcal{R}(\lambda_0) \supset \mathcal{C}(\lambda_0) \setminus X$ and X does not contain $\mathcal{C}(\lambda_0)$ if $\mathcal{R}(\lambda_0)$ is nonempty.*

More precisely, there exists a finite set of angles $E = E(\lambda_0) \subset \text{supp}(\lambda_0)$ such that $f \in \mathcal{C}(\lambda_0)$ is λ_0 -renormalizable if the landing point of $R_f(v, \theta)$ is neither parabolic nor critical for any $(v, \theta) \in E$.

In particular, for a one-dimensional algebraic set $Y \subset \text{Poly}(T_0)$ such that $Y \cap \mathcal{R}(\lambda_0) \neq \emptyset$,

$$(\mathcal{C}(\lambda_0) \setminus \mathcal{R}(\lambda_0)) \cap Y$$

is finite, that is, there exist only finitely many $f \in \mathcal{C}(\lambda_0) \cap Y$ such that f is not λ_0 -renormalizable.

Note that $\mathcal{C}(\lambda_0)$ might be contained in X , and also notice that λ_0 is primitive if and only if E (hence X) can be taken as the empty set.

6.4. Combinatorial tuning. Let $T_0 = (|T_0|, \sigma_0, \delta_0)$ be a mapping schema. Let λ_0 be a T_0 -invariant rational lamination and fix an internal angle system $(\alpha_w : \bar{w} \rightarrow \mathbb{R}/\mathbb{Z})_{w \in |T(\lambda_0)|}$.

Definition (Combinatorial straightening). Let λ be a T_0 -invariant rational lamination containing λ_0 . the *combinatorial straightening of λ with respect to λ_0* is a $T(\lambda_0)$ -invariant rational lamination $\lambda' = (\lambda'_w)_{w \in |T(\lambda_0)|}$ such that $(w, \theta) \sim_{\lambda'_w} (w, \theta')$ if and only if there exist $t \in \alpha_w^{-1}(\theta)$ and $s \in \alpha_w^{-1}(\theta')$ such that t and s are λ -equivalent.

Combinatorial tuning is the inverse operation of combinatorial straightening.

Theorem 6.11 (Combinatorial tuning). *Let λ_0 be a T_0 -invariant rational lamination and let λ' be a $T(\lambda_0)$ -invariant rational lamination. Then there exists a T_0 -invariant rational lamination whose combinatorial straightening with respect to λ_0 is λ' . Moreover,*

- (i) if λ_0 and λ' are hyperbolic, then λ is hyperbolic.
- (ii) If λ_0 and λ' are post-critically finite, then λ is post-critically finite.
- (iii) If λ_0 is post-critically finite and λ' is Misiurewicz, then λ is Misiurewicz.
- (iv) If the rational lamination of $f \in \mathcal{R}(\lambda_0)$ is λ , then the rational lamination of $\chi_{\lambda_0}(f)$ is λ' .

By using the combinatorial tuning, we can actually do “tuning” in most cases of post-critically finite dynamics [IK08, Theorem 6.2]:

Theorem 6.12 (Post-critically finite tuning). *Let λ_0 be a rational lamination over a mapping schema T_0 such that $\mathcal{R}(\lambda_0) \neq \emptyset$.*

Then there exists a codimension one algebraic set Y such that if $P \in \mathcal{C}(T(\lambda_0)) \setminus Y$ is post-critically finite, then there exists $f \in \mathcal{R}(\lambda_0)$ such that $\chi_{\lambda_0}(f) = P$.

Furthermore, if λ_0 is post-critically finite, then such f is unique.

The algebraic set Y in the theorem is defined in a similar way as in Theorem 6.10. In particular, it is empty when λ_0 is primitive. We call f the *tuning* of λ_0 and P , or when $f_0 \in \text{Poly}(T_0)$ satisfies $\lambda_{f_0} = \lambda_0$, we also say f is the *tuning* of f_0 and P . If λ_0 is post-critically finite, then such f is also post-critically finite.

Lemma 6.13. *Let λ_0 be a rational lamination over a mapping schema T_0 and let λ be a $T(\lambda_0)$ -invariant rational lamination.*

If λ is primitive and all periods of periodic λ -unlinked classes are sufficiently large, then the combinatorial tuning λ_1 of λ_0 and λ is also primitive.

Proof. Assume λ_1 is not primitive, that is, there exist some λ_1 -unlinked class L_1 and L_2 and λ_1 -class A such that $\overline{L_j} \cap A \neq \emptyset$ for $j = 1, 2$.

If L_1 and L_2 lie in the same λ_0 -unlinked class M , then there exists some $n \geq 0$ such that $v = m_{T_0}^n(M) \in |T(\lambda_0)|$ and $m_{T_0}^n : M \rightarrow v$ is a cyclic order preserving bijection. This implies that $L'_1 = \alpha_v(m_{T_0}^n(L_1))$ and $L'_2 = \alpha_v(m_{T_0}^n(L_2))$ are λ -unlinked class and there exists a λ -class B intersecting both $\overline{L'_1}$ and $\overline{L'_2}$. Therefore λ is not primitive, that is a contradiction.

Let M_j be the λ_0 -unlinked class containing L_j . We have proved $M_1 \neq M_2$. We also have $\overline{M_j} \cap A \neq \emptyset$. Let $B \subset A$ be the λ_0 -class such that $\overline{M_1} \cap B \neq \emptyset$. Then there exists λ_0 -unlinked class $M_3 \neq M_1$ such that $\overline{M_3} \cap B$ are non-empty. Since the eventual periods of A and B are the same, there are only finite possibility for the eventual period of A by Lemma 6.5.

Therefore, if all periods of periodic λ -unlinked classes are sufficiently large, all periodic angles in the closures of λ -unlinked classes have periods greater than that of such A by Lemma 6.6. Hence the above argument shows that there are no such triple (L_1, L_2, A) , so λ is primitive. $\square$

7. CONTINUITY OF STRAIGHTENING MAPS

Now we give some sufficient condition for straightening maps to be continuous at some $f \in \mathcal{R}(\lambda_0)$.

The argument on continuity of straightening maps by Douady and Hubbard (Theorem 3.2, Lemma 3.3) can be applied to our case. In particular, we have the following:

Lemma 7.1. *Let λ_0 be a rational lamination over T_0 . Assume f_n converges to f in $\mathcal{R}(\lambda_0)$ and $\chi_{\lambda_0}(f_n)$ converges to $h \in \mathcal{C}(T(\lambda_0))$. Then there exist some $K \geq 1$ independent of n and a K -quasiconformal hybrid conjugacy ψ_n between a λ_0 -renormalization of f_n and $\chi_{\lambda_0}(f_n)$ such that ψ_n converges to a K -quasiconformal conjugacy ψ between λ_0 -renormalization of f_n and h by passing to a subsequence. In particular, h and $\chi_{\lambda_0}(f)$ are quasiconformally equivalent.*

Theorem 7.2. *Let λ_0 be a rational lamination over T_0 . If $f \in \mathcal{R}(\lambda_0)$ is quasiconformally rigid, then χ_{λ_0} is continuous at f .*

We say $f \in \text{Poly}(T_0)$ is *quasiconformally rigid* if $g \in \text{Poly}(T_0)$ is quasiconformally conjugate to f , g is affinely conjugate to f .

Proof. Assume f_n converges to f in $\mathcal{R}(\lambda_0)$. Then by the lemma above, we may assume that $\chi_{\lambda_0}(f_n)$ converges to h , which is quasiconformally conjugate to $\chi_{\lambda_0}(f)$. Hence it follows that there exists $\tilde{f} \in \mathcal{R}(\lambda_0)$ such that $\chi_{\lambda_0}(\tilde{f}) = h$ and $\tilde{f}$ is quasiconformally conjugate to f [IK08, Lemma 8.6].

By assumption, we have $\tilde{f} = f$ and $h = \chi_{\lambda_0}(f)$. $\square$

Similarly, we have some partial continuity of $\chi_{\lambda_0}^{-1}$. Observe that when λ_0 is post-critically finite, $\chi_{\lambda_0}^{-1} : \chi_{\lambda_0}(\mathcal{R}(\lambda_0)) \rightarrow \mathcal{R}(\lambda_0)$ is well-defined since χ_{λ_0} is injective.

Proposition 7.3. *Under the assumption of Theorem 7.2, assume λ_0 is post-critically finite and there exists a convergent sequence $f_n \rightarrow \tilde{f}$ in $\mathcal{R}(\lambda_0)$ such that $\chi_{\lambda_0}(f_n) \rightarrow \chi_{\lambda_0}(f)$. Then $\tilde{f} = f$.*

Proof. By Lemma 7.1, $\chi_{\lambda_0}(f)$ and $\chi_{\lambda_0}(\tilde{f})$ are quasiconformally conjugate. Hence, by [IK08, Lemma 8.6], there exists some $\hat{f}$ quasiconformally conjugate to f such that $\chi_{\lambda_0}(\hat{f}) = \chi_{\lambda_0}(\tilde{f})$.

Therefore, $f = \tilde{f}$ by assumption and Theorem 6.7. $\square$

8. PARABOLIC BIFURCATION

In this section and the next section, we study perturbations in the connectedness locus and see when a polynomial having a polynomial-like restriction satisfies the perturbation condition in Theorem 5.1 for sufficiently many repelling periodic point.

In this section, we study parabolic bifurcations and give a sufficient condition to have nice perturbations. The successive section is devoted to the study of Misiurewicz bifurcations to find parabolic polynomials satisfying this sufficient condition.

Definition. We say a polynomial f of degree $d \geq 3$ satisfies (C1) if the following hold;

- (C1-a) 0 is a non-degenerate 1-parabolic periodic point of period p for f ;
- (C1-b) there exists a quadratic-like restriction $f^p : V' \rightarrow V$ of f^p containing 0 hybrid equivalent to $z + z^2$;
- (C1-c) let $\omega \in V'$ be the critical point of this quadratic-like restriction. There exists another critical point ω' for f and $N > 0$ such that $f^n(\omega') \notin K(f^p; V', V)$ for $n < N$ and $f^p(\omega) = f^N(\omega') \in K(f^p; V', V)$.

We say f satisfies (C2) if it satisfies (C1) and

- (C2-a) every critical point other than ω and ω' is preperiodic;
- (C2-b) the rational lamination λ_f of f , which is post-critically finite by (C2-a), is primitive.

Remark 8.1. The condition (C2-a) is just to obtain an analytic subset in the parameter space where a desired bifurcation occurs with keeping other dynamical properties. Therefore, for example, we can relax it to admit critical points in bounded attracting basins. In this case, we can use the analytic dependence of the dynamics in the hyperbolic component [MP92] to get such an analytic subset.

The following condition implies that we have nice perturbations to apply Theorem 5.1 (see the proof of Theorem 1).

Definition. We say f satisfies (C3) if it satisfies (C1) and for any repelling periodic point $\alpha \in K(f^p, V', V)$, there exists a convergent double sequence

$$f_{n,m} \xrightarrow{m \rightarrow \infty} f_n \xrightarrow{n \rightarrow \infty} f$$

in $\mathcal{C}(d)$ such that the following hold. Let us denote the continuations of $z = \omega, \omega', \alpha$ for f_n and $f_{n,m}$ by z_n and $z_{n,m}$ (that is, we require that they do not bifurcate under these perturbations) and let

$$\begin{aligned} x_n &= f_n^p(\omega_n), & y_n &= f_n^N(\omega'_n), \\ x_{n,m} &= f_{n,m}^p(\omega_{n,m}), & y_{n,m} &= f_{n,m}^N(\omega'_{n,m}). \end{aligned}$$

(Recall that $\lim x_n = \lim y_n$ by (C1-c).)

- (C3-a) 0 is a periodic point of period p for f_n and $f_{n,m}$. It is non-degenerate and 1-parabolic for f_n ;
- (C3-b) $x_n \neq y_n$ (hence $x_{n,m} \neq y_{n,m}$ for sufficiently large m).
- (C3-c) the other critical orbit relations of f are preserved for $f_{n,m}$ (hence also for f_n), i.e., all critical points do not bifurcate under these perturbations and if $c, c' \in \text{Crit}(f) \setminus \{\omega, \omega'\}$ (possibly $c = c'$) satisfy $f^k(c) = f^{k'}(c')$, then $f_{n,m}^k(c_{n,m}) = f_{n,m}^{k'}(c'_{n,m})$ for any n, m where $c_{n,m}$ and $c'_{n,m}$ are the continuations of c and c' respectively.
- (C3-d) $f_{n,m}^p : V'_{n,m} \rightarrow V_{n,m}$ is a quadratic-like restrictions near 0 and w , converging to a quadratic-like restriction $f_n^p : V'_n \rightarrow V_n$ locally uniformly, and it also converges to $f^p : V' \rightarrow V$ as $n \rightarrow \infty$. (Hence $f_n^p : V'_n \rightarrow V_n$ is hybrid equivalent to $z + z^2$.)
- (C3-e) $x_n, y_n \in \text{int}K(f_n^p; V'_n, V_n)$, and $x_{n,m}, y_{n,m} \in K(f_{n,m}^p; V'_{n,m}, V_{n,m})$.
- (C3-f) $f_{n,m}$ geometrically converges to (f_n, g_n) as $m \rightarrow \infty$ such that $g_n(x_n) = \alpha_n$ and $g'_n(x_n) \neq 0$.

Remark 8.2. For simplicity, we say a polynomial satisfies (C1), (C2), or (C3) if a polynomial affinely conjugate to it does, because we mainly consider the space of monic centered polynomials.

We can also define these conditions for a polynomial over a mapping schema in the same way. Observe that the mapping schema must have non-trivial Fatou critical relation in order to satisfy those conditions.

Here, we prove the following.

Theorem 8.3. *Let λ_0 be a post-critically finite d -invariant rational lamination with non-trivial Fatou critical relation. Assume $f \in \mathcal{R}(\lambda_0)$ satisfies (C2) and $\mathcal{R}(\lambda_f) \subset \mathcal{R}(\lambda_0)$. Then f satisfies (C3) such that $f_n, f_{n,m} \in \mathcal{R}(\lambda_0)$ and $\lambda_{f_n} = \lambda_f$ for any n, m .*

The rest of this section is devoted to prove this theorem. We first study the bifurcation of a quadratic polynomial $Q(z) = z^2 + 1/4 \in \text{Poly}(2)$, which is affinely conjugate to $z + z^2$. Consider a repelling periodic point $\alpha(Q)$ of Q and let θ be the landing angle for $\alpha(Q)$. Let c_m be the landing point of the parameter ray $\mathcal{R}_{\mathcal{M}}(\theta/2^m)$ for the Mandelbrot set and let $Q_m(z) = z^2 + c_m$. Let $\alpha(Q_m)$ be the landing point of the external ray $R_{Q_m}(\theta)$, which is the repelling periodic point and $\alpha(Q_m) \rightarrow \alpha(Q)$ as $m \rightarrow \infty$. The critical point 0 is preperiodic under Q_m because $c_m = Q_m(0)$ is the landing point of $R_{Q_m}(\theta/2^m)$ [DH85], it follows that $Q_m^{m+1}(0) = \alpha(Q_m)$ and Q_m is Misiurewicz.

Lemma 8.4. *There exists some Lavaurs map g_Q such that $Q_m \xrightarrow{\text{geom}} (Q, g_Q)$ with $g_Q(Q(0)) = \alpha(Q)$ and $g'_Q(Q(0)) \neq 0$.*

Proof. Since the Mandelbrot set is locally connected at $1/4$ [Hub93], $c_m \rightarrow 1/4$ as $m \rightarrow \infty$. Furthermore, this convergence is tangential to the positive real axis, hence it follows that a Fatou coordinate Φ_{Q_m} is defined for sufficiently large $m > 0$.

By the continuity of Fatou coordinates, there exists some $k > 0$ such that the landing point of $R_{Q_m}(\theta/2^k)$ is contained in the domain of definition of Φ_{Q_m} for sufficiently large

$m > 0$. We may also assume c_m is also contained in the domain of definition of Φ_{Q_m} because we can extend Φ_{Q_m} by the functional equation $\Phi_{Q_m}(Q_m(z)) = \Phi_{Q_m}(z) + 1$. Since Φ_{Q_m} has a critical point only at the backward orbit of the critical point, Φ_{Q_m} is univalent on a neighborhood of $1/4$ and c_m of a definite size. Hence $Q_m^m(c_m) = Q_m^k \circ \Phi_{Q_m}^{-1}(\Phi_{Q_m}(z) + m - k)$ is well-defined and univalent near $z = c_m$ because Q_m^k is univalent on a neighborhood of $\Phi_{Q_m}^{-1}(\Phi_{Q_m}(z) + m - k)$, which is the landing point of $R_{Q_m}(\theta/2^k)$, of a definite size.

Therefore, $Q_m^k \circ \Phi_{Q_m}^{-1}(\Phi_{Q_m}(z) + m - k)$ converges to a Lavaurs map g_Q as $m \rightarrow \infty$, which is univalent near $1/4$. $\square$

Take a sequence $\{\theta_n\} \subset \mathbb{Q}/\mathbb{Z}$ such that $\theta_n \rightarrow \theta$, and let $\alpha_n(Q_m)$ be the landing point of the external ray $R_{Q_m}(\theta_n)$. Since Q_m is Misiurewicz, $J(Q_m)$ is locally connected and hence $\alpha_n(Q_m) \rightarrow \alpha(Q_m)$ as $n \rightarrow \infty$. Let $y_n(Q_m)$ be the landing point of $R_{Q_m}(\theta_n/2^m)$ and we may assume $y_n(Q) = \lim_{m \rightarrow \infty} y_n(Q_m)$ exists (see Figure 2).




FIGURE 2. The external rays of angle θ , θ_n , $\theta/2^m$, $\theta_n/2^m$ and $\theta/2^{m+1}$ for Q_m and their limits. By construction, $R_{Q_m}(\theta)$ and $R_{Q_m}(\theta_n)$ land at repelling (pre)periodic points $\alpha(Q_m)$ and $\alpha_n(Q_m)$ respectively, and $R_{Q_m}(\theta/2^m)$ and $R_{Q_m}(\theta/2^{m+1})$ land at the critical value and critical point respectively.

Let f satisfy the assumption of the Theorem. By assumption, the reduced mapping schema $T(\lambda_f) = (|T(\lambda_f)|, \sigma, \delta)$ is equal to T_{cap, d_1+1} , the schema of capture type of degree $d_1 + 1$.

Define a polynomial $\tilde{Q}_{n,m}$ over $T(\lambda_f)$ as follows:

$$\tilde{Q}_{n,m}(v_i, z) = \begin{cases} (v_0, Q_m(z)) & \text{if } i = 0, \\ (v_0, z^{d_1} + y_n(Q_m)) & \text{if } i = 1. \end{cases}$$

Then $\tilde{Q}_n = \lim_{m \rightarrow \infty} \tilde{Q}_{n,m}$ and $\tilde{Q} = \lim_{n \rightarrow \infty} \tilde{Q}_n$ exist and satisfy the following:

$$\begin{aligned}\tilde{Q}_n(v_i, z) &= \begin{cases} (v_0, Q(z)) & \text{if } i = 0, \\ (v_0, z^{d_1} + y_n(Q)) & \text{if } i = 1, \end{cases} \\ \tilde{Q}(v_i, z) &= \begin{cases} (v_0, Q(z)) & \text{if } i = 0, \\ (v_0, z^{d_1} + \frac{1}{4}) & \text{if } i = 1. \end{cases}\end{aligned}$$

Since $\tilde{Q}_{n,m}$ is Misiurewicz and λ_f is primitive, there exists $f \in \mathcal{R}(\lambda_f) = \mathcal{C}(\lambda_f)$ such that $\chi_{\lambda_f}(f) = \tilde{Q}_{n,m}$ by Theorem 6.12.

By taking a subsequence, we may assume

$$f_{n,m} \xrightarrow{m \rightarrow \infty} f_n \xrightarrow{n \rightarrow \infty} \hat{f}$$

for some f_n and $\hat{f}$. Since $\mathcal{C}(\lambda_f) = \mathcal{R}(\lambda_f)$ is compact by Theorem 6.9, f_n and $\hat{f}$ are also λ_f -renormalizable.

Let $(f_n^{\ell_i} : U'_{n,v_i} \rightarrow U_{n,v_0})_{i=0,1}$ and $(\hat{f}^{\ell_i} : \hat{U}'_{v_i} \rightarrow \hat{U}_{v_0})_{i=0,1}$ be λ_f -renormalizations of f_n and $\hat{f}$. Since the straightening map is continuous for quadratic-like families, the quadratic-like restrictions $f_n^{\ell_0} : U'_{n,v_0} \rightarrow U_{n,v_0}$ and $\hat{f}^{\ell_0} : \hat{U}'_{v_0} \rightarrow \hat{U}_{v_0}$ are hybrid equivalent to $Q(z) = z^2 + 1/4$. Let ω_n and ω'_n (resp. $\hat{\omega}$ and $\hat{\omega}'$) be the critical points for f_n (resp. $\hat{f}$) lying in U'_{n,v_0} and U'_{n,v_1} (resp. $\hat{U}'_{v_0}$ and $\hat{U}'_{v_1}$) respectively. Note that $\ell_0 = p$ and $\ell_1 = N$.

Lemma 8.5. (i) $f_n^N(\omega'_n) \in \text{int}K_{f_n}(v_0)$ for sufficiently large n .
(ii) $\hat{f}^N(\hat{\omega}') = \hat{f}^p(\hat{\omega})$.

Proof. Since $\mathcal{R}(\lambda_f)$ is compact, we may assume that the hybrid conjugacy $\psi_{n,m}$ between λ_f -renormalization of $f_{n,m}$ and $\tilde{Q}_{n,m} = \chi_{\lambda_f}(f_{n,m})$ are uniformly K -quasiconformal by Lemma 7.1. By passing to a subsequence, we may further assume that

$$\psi_{n,m} \xrightarrow{m \rightarrow \infty} \varphi_n \xrightarrow{n \rightarrow \infty} \hat{\varphi}$$

and they are all K -quasiconformal. Then φ_n conjugates the λ_n -renormalization of f_n to $\tilde{Q}_n$ and $\hat{\varphi}$ conjugates that of $\hat{f}$ to $\tilde{Q}$. Hence it follows that

$$\hat{\varphi}(\hat{f}^N(\hat{\omega}')) = \tilde{Q}(\hat{\varphi}(\hat{\omega}')) = (v_0, 1/4) = \hat{\varphi}(\hat{f}^p(\hat{\omega})),$$

so we have (ii) and $\hat{f}^N(\hat{\omega}') \in \text{int}K_{\hat{f}}(v_0)$. Therefore, (i) also follows by continuity. Note that $K(\tilde{Q}_n, v_0) = K(\tilde{Q}, v_0) = K(Q)$ does not depend on n . $\square$

Lemma 8.6. $\lambda_{f_n} = \lambda_{\hat{f}} = \lambda_f$ for sufficiently large n .

Proof. We use the same notation as the proof of the previous lemma. By the previous lemma, the critical points of $\tilde{Q}_n$ and $\tilde{Q}$ lie in the interior of the filled Julia set. Since $K(\tilde{Q}_n, v_0) = K(\tilde{Q}, v_0) = K(Q)$, it follows that the real laminations of $\tilde{Q}_n$ and $\tilde{Q}$ are trivial. By Theorem 6.11, the rational laminations of f_n and $\hat{f}$ are the combinatorial tuning of λ_f and the trivial rational lamination, which is equal to λ_f itself. $\square$

Lemma 8.7. $\hat{f} = f$.

Proof. The quasiconformal rigidity of $\tilde{Q}$ implies that $\chi_{\lambda_f}(\hat{f}) = \chi_{\lambda_f}(f) = \tilde{Q}$. Therefore, the lemma follows from the injectivity of the straightening map χ_{λ_f} . $\square$

Proof of Theorem 8.3. We have already constructed a convergent double sequence

$$f_{n,m} \xrightarrow{m \rightarrow \infty} f_n \xrightarrow{n \rightarrow \infty} f$$

in $\mathcal{R}(\lambda_f) \subset \mathcal{R}(\lambda_0)$, hence it is enough to check that this satisfies (C3).

The condition (C3-a) holds by changing the coordinate if necessary. The condition (C3-b) follows from the fact that $\varphi_n(x_n) = \tilde{Q}_n(v_0, 0) \neq \varphi_n(y_n) = \tilde{Q}_n(v_1, 0)$ by construction. Since all critical points except ω and ω' lie in the Julia set and preperiodic, their behavior is described in terms of λ_f . Thus (C3-c) follows because $f_{n,m}$ and f_n admits λ_f . The λ_f -renormalizability of $f_{n,m}$ and f_n and Lemma 8.5 imply (C3-d) and (C3-e).

Let g_Q be the Lavaurs map in Lemma 8.4. Then $g_Q(y_n(Q)) = \alpha_n(Q)$ by construction. Since $Q_m^m(w)$ ($0 \leq k \leq m$) is sufficiently close to $K(Q)$ for w sufficiently close to $1/4$, we have

$$f_{n,m}^{mp}(z) = \psi_{n,m}^{-1} \circ \tilde{Q}_{n,m}^m(\psi_{n,m}(z)) \rightarrow \varphi_n^{-1} \circ \tilde{g}_n^Q(\varphi_n(z))$$

for z sufficiently close to x_n , where $\tilde{g}_n^Q(v_0, w) = \lim_{m \rightarrow \infty} \tilde{Q}_{n,m}^m(v_0, w) = (v_0, g_Q(w))$. Therefore, this converges to a Lavaurs map g_n such that $g_n = \varphi_n^{-1} \circ \tilde{g}_n^Q \circ \varphi_n$. Since φ_n is quasiconformal and $g'_Q(Q(0)) \neq 0$, we have $g'_n(x_n) \neq 0$. Moreover,

$$g_n(x_n) = \varphi_n^{-1}(v_0, g_Q(Q(0))) = \varphi_n^{-1}(v_0, \alpha(Q)) = \alpha_n.$$

Therefore, we have proved (C3-f). $\square$

9. MISIUREWICZ BIFURCATION

In this section, we prove the following:

Theorem 9.1. *Let λ_0 be a post-critically finite d -invariant rational lamination with a non-trivial Fatou critical relation and let $f_0 \in \mathcal{R}(\lambda_0)$ be a Misiurewicz λ_0 -renormalizable polynomial.*

Then there exists a polynomial $f \in \mathcal{R}(\lambda_0)$ arbitrarily close to f_0 such that

- (i) *f satisfies (C2).*
- (ii) *$\mathcal{R}(\lambda_f) \subset \mathcal{R}(\lambda_0)$.*

The main difficulties of the proof of this theorem are the following. First, all perturbation must be done inside $\mathcal{R}(\lambda_0)$. In order to do this, we perturb in $\mathcal{C}(T(\lambda_0))$ and use tuning to get nice perturbations. Secondly, tuning is not defined everywhere, nor continuous. What we want to apply the tuning is a parabolic map, but tuning does not have such kind of nice properties for parabolic maps. Therefore, we approximate by Misiurewicz maps with a help of combinatorial continuity by Kiwi [Kiw05] and apply Proposition 7.3 to show that what we construct is close to the original map.

Remark 9.2. Similar to Remark 8.1, what we essentially need is a two-dimensional analytic set passing through f_0 , on which we have two active critical points, and the rest of the dynamics behaves stable.

9.1. Critical portraits and combinatorial continuity. Here we briefly recall the notion of critical portraits and the combinatorial continuity for Misiurewicz maps.

Definition. Let T be a mapping schema. A *critical portrait* over T is a collection of sets $\Theta = \{\Theta_1, \dots, \Theta_m\}$ such that each Θ_j is contained in a fiber $\{v_j\} \times \mathbb{R}/\mathbb{Z}$ for some $v_j \in |T|$ and

- (CP1) for every j , $\#\Theta_j \geq 2$ and $\#m_T(\Theta_j) = 1$;
- (CP2) $\Theta_1, \dots, \Theta_m$ are pairwise unlinked;

$$(CP3) \sum_{j=1}^m (\#\Theta_j - 1) = d - 1.$$

We say a critical portrait Θ is *preperiodic* if all elements in $\Theta_1, \dots, \Theta_m$ are preperiodic by $m_T(v, \theta) = (\sigma(v), m_{\delta(v)}(\theta))$.

For a Misiurewicz polynomial P over T , Θ is a *critical portrait of P* if for each j , there exists a critical point ω_j such that $R_f(v, \theta)$ lands at ω_j for any $(v, \theta) \in \Theta_j$. In this case, Θ is always preperiodic.

We endow the space of all critical portraits over T with the *compact-unlinked topology*, which is generated by the subbasis formed by

$$V_X = \{\Theta = \{\Theta_j\}; X \text{ is unlinked with } \Theta_j \ (\forall j)\}$$

where X is a closed subset of $\{v\} \times \mathbb{R}/\mathbb{Z}$ for some $v \in |T|$.

Let us denote by $\mathcal{A}(T)$ the set of critical portraits over T and let

$$\text{Preper}(\mathcal{A}(T)) = \{\Theta \in \mathcal{A}(T); \text{preperiodic}\}.$$

For each critical portrait Θ , we can naturally associate the *impression* of Θ , which is a set $I(\Theta) \subset \mathcal{C}(T) \cap \mathcal{S}(T)$, where $\mathcal{S}(T)$ is the *shift locus*, i.e., the set of polynomials over T with all critical points escaping [Kiw05]. We can also associate a real lamination λ_Θ . Kiwi proved the following [Kiw05, Theorem 1, Corollary 5.3]:

Theorem 9.3 (Kiwi). *For a critical portrait Θ , either*

- *(aperiodic case) for any $f \in I(\Theta)$, all cycles of f are repelling and λ_Θ coincides with the real lamination of f . Each $\Theta_j \in \Theta$ is contained in a λ_Θ -equivalence class, or*
- *(periodic case) any $f \in I(\Theta)$ contains at least one non-repelling cycle.*

Furthermore if Θ is preperiodic, then Θ is aperiodic and $I(\Theta)$ consists of a unique Misiurewicz polynomial f_Θ over T such that Θ is a critical portrait of f_Θ .

In particular, a map

$$\text{Preper}(\mathcal{A}(T)) \rightarrow \text{Mis}(\text{Poly}(T)), \quad \Theta \mapsto f_\Theta \in \text{Poly}(T),$$

is well-defined and continuous, where $\text{Mis}(\text{Poly}(T)) = \{f \in \text{Poly}(T); \text{Misiurewicz}\}$.

When the Julia set of $f \in \mathcal{C}(T)$ is locally connected, the *real lamination* of f is the equivalence relation on $\mathbb{R}/\mathbb{Z}$ defined by the landing relation of external rays, as in the case of the rational lamination (see [Kiw05] for general case).

Lemma 9.4. *Let $f \in \mathcal{R}(\lambda_0)$ be Misiurewicz. Then there exists a critical portrait $\Theta = \Theta(f) = \{\Theta_1, \dots, \Theta_m\}$ of f such that*

- *each Θ_j is either contained in some $A \in \text{Crit}^P(\lambda_0)$ or the closure of a unique $v = v(j) \in |T(\lambda_0)|$, and*

$$\sum_{\Theta_j \subset \bar{v}} (\#\Theta_j - 1) = \delta(v) - 1$$

for each $v \in |T(\lambda_0)|$.

- *Let*

$$\alpha(\Theta) = \left\{ \{v(j)\} \times \alpha_{v(j)}(\Theta_j); \Theta_j \subset \overline{v(j)} \text{ for some } v(j) \in |T(\lambda_0)| \right\}.$$

Then $\alpha(\Theta)$ is a critical portrait of $g = \chi_{\lambda_0}(f)$, where $\alpha = (\alpha_v : \bar{v} \rightarrow \mathbb{R}/\mathbb{Z})$ is the (previously given) internal angle system of λ_0 .

Proof. Since f is λ_0 -renormalizable, each critical point ω of f is either contained in $K_f(v)$ for a unique $v \in \text{Crit}^F(\lambda_0) = |T(\lambda_0)|$, or the landing point of the external rays of angle in $A \in \text{Crit}^P(\lambda_0)$ and not contained in any $K_f(v)$ for $v \in CO^F(\lambda_0)$. In particular, $A \in \text{Crit}^P(\lambda_0)$ does not intersect $\bar{v}$ for any $v \in CO^F(\lambda_0)$.

Therefore, we can choose a critical portrait $\Theta = \{\Theta_1, \dots, \Theta_m\}$ of f so that Θ_j is either contained in $\bar{v}$ for some $v = v(j) \in \text{Crit}^F(\lambda_0)$, or $A \in \text{Crit}^P(\lambda_0)$.

In the former case, there exists a critical point $\omega_j \in K_f(v(j))$ such that $R_f(\theta)$ lands at ω_j for any $\theta \in \Theta_j$ by definition. Since $f \in \mathcal{R}(\lambda_0)$, it follows that $K_f(v) \cap \text{Crit}(f) = \{\omega_j; \theta_j \subset \bar{v}\}$ and

$$\sum_{\Theta_j \subset \bar{v}} (\#\Theta_j - 1) \leq \sum_{\omega_j \in K_f(v) \cap \text{Crit}(f)} (\deg_{\omega_j}(f)) = \delta(v) - 1.$$

By (CP3) and the fact $\sum_{A \in \text{Crit}(\lambda_0)} (\delta(A) - 1) = d - 1$, the above inequality must be an equality.

Furthermore, by the definition of α , $R_g(v(j), \alpha_{v(j)}(\theta))$ also lands at a critical point of g for $\theta \in \Theta_j \subset \bar{v}$. Hence $\alpha(\Theta)$ is a critical portrait of $\chi_{\lambda_0}(f)$. $\square$

9.2. Perturbation in the target space. To prove Theorem 9.1, we first need to construct some nice perturbations in the target space, i.e., we perturb $P_0 = \chi_{\lambda_0}(f_0)$.

First, we need the theorem on universality of the Mandelbrot set by McMullen [McM00] to find such f . For a mapping schema T , let $T^{\text{per}} = (|T^{\text{per}}|, \sigma, \delta)$ be a sub-schema such that $|T^{\text{per}}| = \{v \in |T|; \text{periodic}\}$. Then we have natural projection $\pi : \text{Poly}(T) \rightarrow \text{Poly}(T^{\text{per}})$. We call $\pi(P)$ the *periodic part of P* for $P \in \text{Poly}(T)$.

Theorem 9.5. *Let T be a mapping schema and consider a holomorphic one-parameter family $(P_\mu)_{\mu \in \Delta}$ in $\text{Poly}(T)$ parameterized by the unit disk Δ . Let $\mathcal{B}$ be the bifurcation locus of the family $(\pi(P_\mu))$ of the periodic parts of (P_μ) . Then either $\mathcal{B}$ is empty or there exists a quasiconformal image $\mathcal{M}'$ of $\mathcal{M}_\delta$ for some $\delta \geq 2$ whose boundary is contained in $\mathcal{B}$ where $\mathcal{M}_\delta$ is the connectedness locus of the unicritical family $\{z^\delta + c; c \in \mathbb{C}\}$ of degree δ .*

More precisely, there exists some $n > 0$ such that for μ in $\mathcal{M}'$, there exist a critical point ω_μ analytically parameterized by μ and a polynomial-like restriction $P_\mu^n : W'_\mu \rightarrow W_\mu$ hybrid equivalent to $z^\delta + c(\mu)$ such that

- $\omega_\mu \in W'_\mu$,
- the local degree of P_μ at ω_μ is equal to δ , and
- $c : \mathcal{M}' \rightarrow \mathcal{M}_\delta$ extends to a quasiconformal map of the plane.

Proof. This is a simple generalization of [McM00, Theorem 1.1, Theorem 4.1]. For $\mu \in \Delta$, let us denote $P_\mu^n(v, z) = (\sigma^n(v), P_{\mu, v}(z))$. If $\mathcal{B}$ is nonempty, then there exists some periodic $v \in |T|$ by σ such that the family $(P_{\mu, v}^p)_{\mu \in \Delta}$ has nonempty bifurcation locus where p is the period of v . Therefore, it contains the quasiconformal image of $\partial \mathcal{M}_\delta$, for some $\delta \geq 2$. Since $\mathcal{B}$ contains the bifurcation locus of $(P_{\mu, v}^p)$, the theorem follows by [McM00, Theorem 4.1]. $\square$

Now consider the family of polynomials over a mapping schema T of total degree $d = \delta(T)$ with all critical points marked (counted with multiplicity);

$$\widehat{\text{Poly}}(T) = \{(P, (v_1, \omega_1), (v_2, \omega_2), \dots, (v_{d-1}, \omega_{d-1})); P \in \text{Poly}(T), \text{Crit}(P) = \{(v_j, \omega_j)\}\}.$$

Lemma 9.6. *Let $(P_0, (v_j, \omega_{0,j})) \in \widehat{\text{Poly}}(T)$ be Misiurewicz. Then there exist $d-1$ polynomials $h_j = h_j(P, \omega_j)$ for $j = 1, \dots, d-1$ such that $(P_0, (v_j, \omega_{0,j}))$ is discrete in*

$$\{h_1(P, \omega_1) = \dots = h_{d-1}(P, \omega_{d-1}) = 0\}.$$

Proof. Since each critical point is preperiodic, there exists a relation $P_{0,v_j}^{n_j}(\omega_j) = P_{0,v_j}^{m_j}(\omega_j)$ for some $n_j \neq m_j$. Let $h_j(P, \omega_j) = P_{v_j}^{n_j}(\omega_j) - P_{v_j}^{m_j}(\omega_j)$. Since the set of Misiurewicz polynomials does not contain any one-dimensional algebraic set, $(P_0, (v_j, \omega_{0,j}))$ is discrete in $\{h_1(P, \omega_1) = \dots = h_{d-1}(P, \omega_{d-1}) = 0\}$. $\square$

Now consider a Misiurewicz polynomial $(P_0, (v_j, \omega_{0,j})) \in \widehat{\text{Poly}}(T(\lambda_0))$ and its neighborhood $\mathcal{U}$. Let Θ_0 be a critical portrait of P_0 . Take a preperiodic critical portrait Θ close to Θ_0 such that

- $\Theta = \{\Theta_1, \dots, \Theta_{d-1}\}$, i.e., $\#\Theta_j = 2$ for each j ;
- there exists some $N' > 1$ such that $m_T(\Theta_1) = m_T^{N'}(\Theta_2)$ and $m_T^{N'-1}(\Theta_2) \not\subset \Theta_1$;
- let p_j be the eventual period of Θ_j by m_T . then $p_1(=p_2), p_3, \dots, p_{d-1}$ are mutually different.

Let $(P_\Theta, (v_j, \omega_{\Theta,j})) \in \widehat{\text{Poly}}(T(\lambda_0))$ be the polynomial having Θ as a critical portrait and the landing point of the external ray of angle $(v_j, \theta) \in \Theta_j$ is (v_j, ω_j) . Observe that there are no multiple critical points i.e., (v_j, ω_j) are mutually different because otherwise both the eventual periods and the preperiods must coincide. If Θ is sufficiently close to Θ_0 , then $P_\Theta \in \mathcal{U}$ by Theorem 9.3. Moreover, we have

$$P_\Theta^{N'}(v_2, \omega_{\Theta,2}) = P_\Theta(v_1, \omega_{\Theta,1}), \quad P_\Theta^{N'-1}(v_2, \omega_{\Theta,2}) \neq (v_1, \omega_{\Theta,1}).$$

In fact, the first equality is trivial and if $P_\Theta^{N'-1}(v_2, \omega_{\Theta,2}) = (v_1, \omega_{\Theta,1})$, then all of the three external rays of angles in $\Theta_1 \cup m_T^{N'-1}(\Theta_2)$ land at $(v_1, \omega_{\Theta,1})$ and are mapped to the same ray $R_{f_\Theta}(\theta)$, where $\{\theta\} = m_T(\Theta_1) = m_T^{N'}(\Theta_2)$. This implies that $(v_1, \omega_{\Theta,1})$ is a multiple critical point, so it is a contradiction.

Take $h_1, \dots, h_{d-1}$ in Lemma 9.6 and consider an algebraic set

$$(3) \quad \mathcal{X} = \{(P, (v_j, \omega_j)) \in \widehat{\text{Poly}}(T(\lambda_0)); h_3(P, \omega_3) = \dots = h_{d-1}(P, \omega_{d-1}) = h(P, \omega_1, \omega_2) = 0\},$$

where $h(P, \omega_1, \omega_2) = P^{N'}(v_2, \omega_2) - P(v_1, \omega_1)$.

Lemma 9.7. $\dim \mathcal{X} = 1$.

Proof. Consider the following algebraic sets:

$$\begin{aligned} \mathcal{X}' &= \{h_3(P, \omega_3) = \dots = h_{d-1}(P, \omega_{d-1}) = 0\}, \\ \mathcal{X}'' &= \{h_1(P, \omega_1) = h_3(P, \omega_3) = \dots = h_{d-1}(P, \omega_{d-1}) = h(P, \omega_1, \omega_2) = 0\}. \end{aligned}$$

Then $\dim \mathcal{X}' = 2$ and $\dim \mathcal{X}'' = 0$. Since $\mathcal{X}'' \subset \mathcal{X} \subset \mathcal{X}'$ and $\dim \mathcal{X}' - \dim \mathcal{X}, \dim \mathcal{X} - \dim \mathcal{X}'' \leq 1$, the dimension of $\mathcal{X}$ is one. $\square$

Observe that since all critical points are simple for P_Θ , the natural projection of $\widehat{\text{Poly}}(T_0) \rightarrow \text{Poly}(T_0)$ is a local isomorphism at P_Θ . Hence we identify them to simplify the notation.

Let $\mathcal{X}_0$ be an irreducible component of $\mathcal{X}$ passing through P_Θ and let $\mathcal{B}$ be the bifurcation locus of the periodic parts in $\mathcal{X}_0$. Then since P_Θ is Misiurewicz and a free critical point ω_1 is contained in the periodic part, $P_\Theta \in \mathcal{B}$. In particular, $\mathcal{B}$ is nonempty. Therefore, by Theorem 9.5, There exists a copy $\mathcal{M}' \subset \mathcal{U} \cap \mathcal{X}_0$ of the Mandelbrot set $\mathcal{M} = \mathcal{M}_2$, for all critical points are simple. Let $\xi : \mathcal{M}' \rightarrow \mathcal{M}$ be the homeomorphism defined by

straightening. Let P_1 be the center of $\mathcal{M}'$, i.e., the quadratic-like restriction $P_1^p : W_1' \rightarrow W_1$ is hybrid equivalent to z^2 (i.e., $\xi(P_1) = z^2$).

Lemma 9.8. *We can take $\mathcal{M}'$ so that P_1 (equivalently, λ_{P_1}) is primitive and p is arbitrarily large.*

Proof. If P_1 is not primitive, then take a small copy $\mathcal{M}'' \subset \mathcal{M}'$ which corresponds to a primitive copy of sufficiently high period in $\mathcal{M}$. Then the rational lamination of the center $P_2 \in \mathcal{M}''$ is the combinatorial tuning of λ_{P_1} and a primitive rational lamination over $T(\lambda_{P_1}) = T_{\text{cap}}$. Hence P_2 is primitive by Lemma 6.13.

Therefore, the lemma is obtained by replacing $\mathcal{M}'$ by $\mathcal{M}''$. $\square$

Therefore, we have proved the following.

Lemma 9.9. *Let P_0 be a Misiurewicz polynomial over a mapping schema T . For any neighborhood $\mathcal{U}$ of P_0 , there exist a one-dimensional algebraic subset $\mathcal{X}_0 \subset \widehat{\text{Poly}}(T)$ and a small copy of the Mandelbrot set $\mathcal{M}' \subset \mathcal{X}_0 \cap \mathcal{U}$ such that*

- (i) $\mathcal{X}_0$ is a irreducible component of the algebraic set $\mathcal{X}$ defined by the formula (3) for some polynomials $h_3, \dots, h_{d-1}$ and h . In particular, there is essentially only one free critical point on $\mathcal{X}_0$.
- (ii) for any $P \in \mathcal{M}'$, there exists a quadratic-like restriction $P^{p'} : W_P' \rightarrow W_P$ hybrid equivalent to $Q = \xi(P)$ such that the map $\xi : \mathcal{M}' \rightarrow \mathcal{M}$ is a homeomorphism. The period p (depending $\mathcal{M}'$) can be taken arbitrarily large.
- (iii) Let P_1 be the center of $\mathcal{M}'$, i.e., $\xi(P_1) = z^2$. Then λ_{P_1} is primitive.

9.3. Proof of Theorem 9.1. Let λ_0 be a post-critically finite d -invariant rational lamination and let $f_0 \in \mathcal{R}(\lambda_0)$ be Misiurewicz. Take a small neighborhood $\mathcal{V}$ of f_0 . We may assume $\mathcal{V} \cap \mathcal{C}(\lambda_0) \subset \mathcal{R}(\lambda_0)$ by Lemma 6.10. Take a neighborhood $\mathcal{U}$ of $P_0 = \chi_{\lambda_0}(f_0)$ sufficiently small such that

- the codimension one algebraic set in Theorem 6.12 does not intersect $\mathcal{U}$. Therefore, for any post-critically finite $P \in \mathcal{U}$, there exists a unique f such that $\chi_{\lambda_0}(f) = P$, and
- $\chi_{\lambda_0}^{-1} : \text{Mis}(\text{Poly}(T(\lambda_0))) \cap \mathcal{U} \rightarrow \text{Mis}(\text{Poly}(d)) \cap \mathcal{R}(\lambda_0)$ is a homeomorphism into its image and the closure of the image is contained in $\mathcal{V}$.

The existence of such a neighborhood $\mathcal{U}$ is guaranteed by Theorem 7.2 and Proposition 7.3.

Take a sequence of Misiurewicz polynomials $Q_n \in \partial \mathcal{M}$ ($n \geq 2$) such that $Q_n \rightarrow Q_0(z) = z^2 + 1/4$ and let $P_n = \xi^{-1}(Q_n)$. (Recall that P_1 is the center of $\mathcal{M}'$.) Then P_n is also Misiurewicz for $n \geq 2$. Let $f_n = \chi_{\lambda_0}^{-1}(P_n) \in \mathcal{V}$ for $n \geq 1$.

Let $\lambda = \lambda_{f_1}$ be the combinatorial tuning of λ_0 and λ_{P_1} . Since we may assume the period p of quadratic-like renormalization of P_1 arbitrarily large, λ is also primitive by Lemma 6.13. Therefore, $\mathcal{C}(\lambda) = \mathcal{R}(\lambda)$ is compact.

As in Lemma 9.9, there exists a one-dimensional algebraic subset

$$\mathcal{Y} = \{(f, \omega_1, \dots, \omega_{d-1}) \in \widehat{\text{Poly}}(d); \hat{h}_3(\omega_3) = \dots = \hat{h}_{d-1}(\omega_{d-1}) = \hat{h}(f, \omega_1, \omega_2) = 0\}$$

containing all f_n . Since $\mathcal{R}(\lambda)$ is compact, we may assume that f_n converges to some $f \in \mathcal{R}(\lambda)$. Then

$$f \in \mathcal{C}(\lambda) \cap \mathcal{V} \subset \mathcal{C}(\lambda_0) \cap \mathcal{V} = \mathcal{R}(\lambda_0) \cap \mathcal{V}.$$

Namely, f is λ_0 -renormalizable and close to f_0 .

Since $\mathcal{V}$ is closed, f (more precisely, $(f, \omega_1, \dots, \omega_{d-1})$) also lies in $\mathcal{V}$. Take $w_j \in |T(\lambda_0)|$ such that $\omega_j \in K_f(w_j)$ for $j = 1, 2$ and let

$$N = \sum_{j=1}^{N'-1} \ell_{\sigma^j(w_2)}, \quad p = \sum_{j=1}^{p'-1} \ell_{\sigma^j(w_1)},$$

and $N'' = N - (p - \ell_{w_1})$. Then it follows that $f^{\ell_{w_1}}(\omega_1) = f^{N''}(\omega_2)$ (recall that $T(\lambda_0) \cong T_{\text{cap}}$ and $|T_{\text{cap}}| = \{v_1, v_2\}$). By this and Theorem 3.2, we have $\chi_\lambda(f) = \tilde{Q}_0$, where

$$\tilde{Q}(0)(v_j, z) = \begin{cases} (v_0, Q_0(z)) & \text{when } j = 0, \\ (v_0, z^2 + Q_0(0)) & \text{when } j = 1. \end{cases}$$

It is easy to check that f satisfies (C2) (note that p above is different from that in (C1)). $\square$

10. DISCONTINUITY

Now we give a proof of the main theorem:

Proof of Main Theorem. First, observe that there always exists a Misiurewicz polynomial $f_0 \in \mathcal{R}(\lambda_0)$ assuming that $\mathcal{R}(\lambda_0)$ is nonempty, by Theorem 9.3 and Theorem 6.12.

Assume that χ_{λ_0} is continuous on $\mathcal{V} \cap \mathcal{R}(\lambda_0)$ for a neighborhood $\mathcal{V} \in \text{Poly}(T_0)$ of f_0 .

By Theorem 9.1, there exists $f_1 \in \mathcal{V} \cap \mathcal{R}(\lambda_0)$ satisfying (C3). In the following, we use the notations in (C3) like ω , ω' , p , N , V and V' for f_1 . Let $w_0, w_1 \in |T(\lambda_0)|$ satisfy $\omega \in K_{f_1}(w_0)$ and $\omega' \in K_{f_1}(w_1)$. Let s' be the period of w_0 by σ , and s be the period of $K_{f_1}(w_0)$, in other words,

$$s = \sum_{n=0}^{t-1} \ell_{\sigma^n(w_0)}.$$

Similarly, define N' and p' by

$$\sum_{n=0}^{N'-1} \ell_{\sigma^n(w_1)} = N, \quad \sum_{n=0}^{p'-1} \ell_{\sigma^n(w_0)} = p.$$

Observe that $K(f_1^p; V', V) \subset K_{f_1}(w_0)$. In particular, s divides s' and p divides p' .

By shrinking $\mathcal{V}$ if necessary, we may assume any $f \in \mathcal{V}$ has a polynomial-like restriction $g_f = (f^{\ell_w} : U'_{g,w} \rightarrow U_{g,\sigma(w)})_{w \in |T(\lambda_0)|}$ over $T(\lambda_0)$ such that

- (i) g_f is a λ_0 -renormalization when $f \in \mathcal{R}(\lambda_0)$.
- (ii) $(f^s : U''_{f,w_0} \rightarrow U_{f,w_0})_{f \in \mathcal{V}}$ forms an AFPL, where U''_{f,w_0} is the component of $f^{-s}(U_{f,w_0})$ containing $K(g_f, w_0)$.

It follows by Theorem 9.1 that $K(f_1^p; V', V) \subset K_f(w_0)$ for $Qf \in \mathcal{R}(\lambda_0) \cap \mathcal{V}$. Observe that by definition, $g_f^{s'} = f^s$ and $g_f^{p'} = f^p$ on $K_f(w_0)$, and $g_f^{N'} = f^N$ on $K_f(w_1)$.

By taking a finite branched cover of $\mathcal{V}$ if necessary, we may assume there exist analytic parameterizations of critical points $\omega(f)$ and $\omega'(f)$ such that $\omega(f_1) = \omega$ and $\omega'(f_1) = \omega'$. For $f \in \mathcal{R}(\lambda_0) \cap \mathcal{V}$, let $P_f = \chi_{\lambda_0}(f) \in \mathcal{C}(T(\lambda_0))$ and $\psi_f = (\psi_{f,w})_{w \in |T(\lambda_0)|}$ be a hybrid conjugacy between g_f and P_f (we can take such a hybrid conjugacy ψ_f by shrinking $U_{f,w}$ if necessary). Let

$$\omega(P_f) = (w_0, \psi_{f,w_0}(\omega(f))), \quad \omega'(P_f) = (w_1, \psi_{f,w_1}(\omega'(f)))$$

be the critical points corresponding to $\omega(f)$ and $\omega'(f)$ respectively for P_f . Let $x(f) = f^p(\omega(f))$ and $y(f) = f^N(\omega'(f))$ and define $x(P_f)$ and $y(P_f)$ by

$$(w_0, x(P_f)) = P_f^{s'}(\omega(P_f)), \quad (w_0, y(P_f)) = P_f^{N'}(\omega'(P_f)).$$

Observe that $x(P_f), y(P_f) \in K(P_f, w_0)$. Now consider an AFPL2MP

$$\mathbf{h} = (f^s : U_f'' \rightarrow U_f, x(f), y(f))_{f \in \mathcal{V}}.$$

Then the straightening map $\chi_{\mathbf{h}}$ for $\mathbf{h}$ satisfies

$$\chi_{\mathbf{h}}(f) = (\hat{P}_f, \psi_{f, w_0}(f^P(\omega(f))), \psi_{f, w_0}(f^N(\omega'(f)))) = (\hat{P}_f, x(P_f), y(P_f))$$

where $P_f^s(w_0, z) = (w_0, \hat{P}_f(z))$. Since χ_{λ_0} is continuous on $\mathcal{N}$, $\chi_{\mathbf{h}}$ is also continuous.

Consider a repelling periodic point $\alpha = \alpha(f)$ in the filled Julia set $K(f_1^P; V', V)$ of the quadratic-like restriction $f_1^P : V' \rightarrow V$. Then we can take $f_{n,m}, f_n \in \mathcal{R}(\lambda_0) \cap \mathcal{V}$ satisfying the conditions in (C3). Therefore, we can apply Theorem 5.1, namely, we have

$$(4) \quad |\text{mult}_{f_1}(\alpha)| = |\text{mult}_{P_{f_1}}(\psi_{f_1, w_0}(\alpha))|.$$

Observe that ψ_{f_1} is also a hybrid conjugacy from $f_1^P : V' \rightarrow V$ to a quadratic-like restriction of $P_{f_1}^P$. Since (4) holds for any repelling periodic point $\alpha \in K(f_1^P; V', V)$, it follows that $\psi_{f_1}|_V$ preserves multipliers. Therefore, by Theorem 2.2, $\tilde{P}_{f_1}$ and f_1^P are conjugate by an irreducible holomorphic correspondence, where $\tilde{P}_{f_1}$ is defined by $P_{f_1}^P(w_0, z) = (w_0, \tilde{P}_{f_1}(z))$. In particular, $\deg \tilde{P}_{f_1} = \deg f_1^P$. However, since λ_0 is nontrivial, we have $\deg P_{f_1, w} < \deg f_1 \leq \deg f_1^{\ell_w}$ for all w , so $\deg \tilde{P}_{f_1} < (\deg f_1)^P$, that is a contradiction. Therefore, χ_{λ_0} is not continuous on $\mathcal{V}$. $\square$

Remark 10.1. More precisely, we have proved the following: for any repelling periodic point $\alpha \in K(f_1^P; V', V)$ such that (4) does not hold (such a repelling periodic point always exists), there exists a double sequence $f_{n,m} \rightarrow f_n \rightarrow f_1$ satisfying the conditions in (C3) such that

$$\lim_{m \rightarrow \infty} \chi_{\lambda_0}(f_{n,m}) \neq \chi_{\lambda_0}(f_n)$$

for sufficiently large n , because $\chi_{\lambda_0}(f_n) \rightarrow \chi_{\lambda_0}(f_1)$ by the quasiconformal rigidity of f_1 .

11. THE CASE OF RATIONAL AND TRANSCENDENTAL ENTIRE MAPS

We do not know very much how rich the dynamics in a renormalizable set is for families of rational maps and transcendental entire maps. However, since the target space of a straightening map is a family of polynomials over a mapping schema, we can apply the same argument to obtain the following:

Theorem 11.1. *Let $(f_\mu)_{\mu \in \Lambda}$ be an analytic family of rational maps of degree $d \geq 3$. Assume there exists an (externally marked) AFPL $\mathbf{g} = (g_\mu = (f_\mu^{\ell_v} : U_v \rightarrow U_{\sigma(v)})_{v \in |T|})_{\mu \in \Lambda}$ over a mapping schema $T = (|T|, \sigma, \delta)$ having a non-trivial critical relation. Let $\chi : \mathcal{C}(\mathbf{g}) \rightarrow \mathcal{C}(T)$ be the straightening map for $\mathbf{g}$. For a Misiurewicz map $P_0 \in \mathcal{C}(T)$, assume there exist a neighborhood $\mathcal{U}$ of P_0 and a map $s : \mathcal{U} \cap \mathcal{C}(T) \rightarrow \mathcal{C}(\mathbf{g})$ such that $\chi \circ s$ is the identity. Then s is not continuous, except when (f_μ) is affinely conjugate to a family of polynomials and $\delta(v) = d$ for all $v \in |T|$.*

In particular, there is no homeomorphic restriction of χ onto $\mathcal{U} \cap \mathcal{C}(T)$.

An (externally marked) AFPL over a mapping schema, its connectedness locus and its straightening map in the same way.

Proof. Let $P_1 \in \mathcal{C}(T) \cap \mathcal{U}$ satisfy (C3) and let $f_1 = s(P_1) \in \mathcal{C}(\mathbf{g})$. Then the same argument as Theorem 1 can be applied to s to show the discontinuity. $\square$

Theorem 11.2. *Let $(f_\mu)_{\mu \in \Lambda}$ be an analytic family of transcendental entire maps of degree $d \geq 3$. Assume there exists an (externally marked) AFPL $\mathbf{g} = (g_\mu = f_\mu^{\ell_v} : U_v \rightarrow U_{\sigma(v)})_{v \in |T|}$ over a mapping schema $T = (|T|, \sigma, \delta)$ having a non-trivial critical relation. Let $\chi : \mathcal{C}(\mathbf{g}) \rightarrow \mathcal{C}(T)$ be the straightening map for $\mathbf{g}$. Let $P_0 \in \mathcal{C}(T)$ be Misiurewicz and assume there exist a neighborhood $\mathcal{U}$ of P_0 and a continuous map $s : \mathcal{U} \cap \mathcal{C}(T) \rightarrow \mathcal{C}(\mathbf{g})$ such that $\chi \circ s$ is the identity.*

Then there exist some $P_1 \in \mathcal{U} \cap \mathcal{C}(T)$ satisfying (C3), a polynomial g , φ_1 and a transcendental entire map φ_2 such that

$$P_1 \circ \varphi_1 = \varphi_1 \circ g, \quad f_1 \circ \varphi_2 = \varphi_2 \circ g,$$

where $f_1 = s(P_1)$.

The proof is the same as Theorem 11.1. The only difference is that we cannot get a contradiction after applying Theorem 2.6, because the degree of a transcendental entire map is infinite and we cannot exclude the case in the conclusion. Note that it follows that g and φ_1 are polynomials by comparing the growth at the infinity (see [Ino08]).

REFERENCES

- [BH01] X. Buff and C. Henriksen, *Julia sets in parameter spaces*, Comm. Math. Phys. **220** (2001), no. 2, 333–375. MR1844629 (2002d:37075)
- [DF08] Romain Dujardin and Charles Favre, *Distribution of rational maps with a preperiodic critical point*, Amer. J. Math. **130** (2008), no. 4, 979–1032. MR2427006
- [DH85] A. Douady and J. H. Hubbard, *Étude dynamique des polynômes complexes. Partie I & II*, Publications Mathématiques d’Orsay [Mathematical Publications of Orsay], vol. 84–85, Université de Paris-Sud, Département de Mathématiques, Orsay, 1984–85, With the collaboration of P. Lavaurs, Tan Lei and P. Sentenac. MR762431 (87f:58072a), MR812271 (87f:58072b)
- [DH85] Adrien Douady and John Hamal Hubbard, *On the dynamics of polynomial-like mappings*, Ann. Sci. École Norm. Sup. (4) **18** (1985), no. 2, 287–343. MR816367 (87f:58083)
- [Dou94] Adrien Douady, *Does a Julia set depend continuously on the polynomial?*, Complex dynamical systems (Cincinnati, OH, 1994), Proc. Sympos. Appl. Math., vol. 49, Amer. Math. Soc., Providence, RI, 1994, pp. 91–138. MR1315535
- [DSZ97] Adrien Douady, Pierrette Sentenac, and Michel Zinsmeister, *Implosion parabolique et dimension de Hausdorff*, C. R. Acad. Sci. Paris Sér. I Math. **325** (1997), no. 7, 765–772. MR1483715 (98i:58195)
- [Eps] Adam Epstein, *manuscript*.
- [Hai00] Peter Haïssinsky, *Modulation dans l’ensemble de Mandelbrot*, The Mandelbrot set, theme and variations, London Math. Soc. Lecture Note Ser., vol. 274, Cambridge Univ. Press, Cambridge, 2000, pp. 37–65. MR1765084 (2002c:37067)
- [Hub93] J. H. Hubbard, *Local connectivity of Julia sets and bifurcation loci: three theorems of J.-C. Yoccoz*, Topological methods in modern mathematics (Stony Brook, NY, 1991), Publish or Perish, Houston, TX, 1993, pp. 467–511. MR1215974 (94c:58172)
- [IK08] Hiroyuki Inou and Jan Kiwi, *Combinatorics and topology of straightening maps i: compactness and bijectivity*, arXiv.org:0809.1262 (2008).
- [Ino06] Hiroyuki Inou, *Homeomorphisms from Julia sets into connectedness loci*, Ergodic Theory Dynam. Systems **26** (2006), no. 4, 1087–1113. MR2246593 (2007f:37070)
- [Ino08] ———, *Extending local analytic conjugacies*, to appear in Trans. Amer. Math. Soc. (2008).
- [Kiw01] Jan Kiwi, *Rational laminations of complex polynomials*, Laminations and foliations in dynamics, geometry and topology (Stony Brook, NY, 1998), Contemp. Math., vol. 269, Amer. Math. Soc., Providence, RI, 2001, pp. 111–154. MR1810538 (2002e:37063)
- [Kiw05] ———, *Combinatorial continuity in complex polynomial dynamics*, Proc. London Math. Soc. (3) **91** (2005), no. 1, 215–248. MR2149856 (2006e:37078)
- [McM00] Curtis T. McMullen, *The Mandelbrot set is universal*, The Mandelbrot set, theme and variations, London Math. Soc. Lecture Note Ser., vol. 274, Cambridge Univ. Press, Cambridge, 2000, pp. 1–17. MR1765082 (2002f:37081)
- [Mil92] John Milnor, *Remarks on iterated cubic maps*, Experiment. Math. **1** (1992), no. 1, 5–24. MR1181083 (94c:58096)

- [Mil00] ———, *Local connectivity of Julia sets: expository lectures*, The Mandelbrot set, theme and variations, London Math. Soc. Lecture Note Ser., vol. 274, Cambridge Univ. Press, Cambridge, 2000, pp. 67–116. MR1765085 (2001b:37073)
- [MP92] John W. Milnor and Alfredo Poirier, *Hyperbolic components in spaces of polynomial maps*, Tech. Report Stony Brook IMS 1992/3, February 1992.
- [Pra96] Eduardo A. Prado, *Teichmüller distance for some polynomial-like maps*, Tech. Report Stony Brook IMS 1996/2. Dynamical Systems 1/15/96, January 1996.
- [PU99] Feliks Przytycki and Mariusz Urbański, *Rigidity of tame rational functions*, Bull. Polish Acad. Sci. Math. **47** (1999), no. 2, 163–182. MR1686679 (2000i:37065)
- [Sch04] Dierk Schleicher, *On fibers and local connectivity of Mandelbrot and Multibrot sets*, Fractal geometry and applications: a jubilee of Benoît Mandelbrot. Part 1, Proc. Sympos. Pure Math., vol. 72, Amer. Math. Soc., Providence, RI, 2004, pp. 477–517. MR2112117 (2006b:37088)
- [Shi98] Mitsuhiro Shishikura, *The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets*, Ann. of Math. (2) **147** (1998), no. 2, 225–267. MR1626737 (2000f:37056)
- [Shi00] ———, *Bifurcation of parabolic fixed points*, The Mandelbrot set, theme and variations, London Math. Soc. Lecture Note Ser., vol. 274, Cambridge Univ. Press, Cambridge, 2000, pp. 325–363. MR1765097 (2001k:37073)
- [Sul87] Dennis Sullivan, *Quasiconformal homeomorphisms in dynamics, topology, and geometry*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986) (Providence, RI), Amer. Math. Soc., 1987, pp. 1216–1228. MR934326 (90a:58160)
- [Thu85] William Thurston, *On the combinatorics of iterated rational maps*, manuscript (1985).
- [Urb97] Mariusz Urbański, *Geometry and ergodic theory of conformal non-recurrent dynamics*, Ergodic Theory Dynam. Systems **17** (1997), no. 6, 1449–1476. MR1488329 (99j:58178)

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN